

**SUPPLEMENTAL LABELING
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON**

ORTHENE[®] 97
EPA REG. NO. 59639-91-5481
EPA SLN NO. OR-000020
Rev. 3

This label valid until December 31, 2012 or until otherwise amended, withdrawn, cancelled or suspended.

This label must be in the possession of the user at the time of pesticide application.
Follow all other applicable directions, restrictions and precautions on the EPA-registered label.

Active Ingredient	By Wt.
Acephate (O,S-Dimethyl acetylphosphoramidothioate)	97%
Other Ingredients	3%
TOTAL	100%

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

See below for additional precautionary statements.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Wash hands thoroughly after handling. Do not allow children or pets to come into contact with treated areas until sprays have dried.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash Hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into the clean clothing.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to birds. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Cover or soil-incorporate spills.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls, waterproof gloves, shoes plus socks and chemical-resistance headgear for overhead exposure.

Crop	Insects*	Amount <i>Orthene</i> 97 Per Acre	Use Directions/Restrictions	Days to Harvest
Cranberries	Blackheaded fireworm Root Weevils	1.0 lb. (.97 lb Acephate)	<p>Apply in water by air, ground or with sprinklers (chemigation).</p> <p>Use a minimum of 2 gal. spray per acre by air. Use sufficient water to give thorough coverage with ground or sprinkler equipment.</p> <p>For Root Weevils, make application night after peak adult emergence and prior to egg laying.</p> <p>If applied through an irrigation system, refer to "Sprinkler Application to Cranberries" below.¹</p> <p>Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.</p> <p>If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.</p>	>75
<p>¹ Sprinkler Irrigation Application to Cranberries: This product may only be applied through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, travelers, big gun, solid set or had move. Do not apply this product through any other type of irrigation system.</p>				

*Pheromone traps are available to monitor flight activity of the Blackheaded Fireworm moth stage and are helpful in predicting egg hatch and early larval stages. Frequent scouting is also advised. Consult local Extension Service personnel and/or Consultants for information on application timings based on trap catches and other Integrated Pest Management (IPM) procedures.

RESTRICTIONS

1. Do not tank mix *Orthene* 97 with other organophosphate insecticides when making application post-bloom.
2. Do not apply from start of bloom until all berries set.
3. Do not apply less than 75 days before harvest.
4. Apply no more than 1.0 lb of Acephate per acre per crop cycle.

FIFRA § 24(c) Registrant: Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596-8025

Distributed By: AMVAC Chemical Corporation
4100 E. Washington Blvd.
Los Angeles, CA 90023
www.Amvac-Chemical.com

Orthene is a registered trade mark of OMS Investments, Inc.

U.S. Pat. 6,013,272