

Supplemental Label



Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596

SUPPLEMENTAL LABEL FOR SPECIAL LOCAL NEED REGISTRATION

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF OREGON

COBRA[®] Herbicide

FOR WEED CONTROL IN NON-BEARING STRAWBERRIES

EPA Reg. No. 59639-34

EPA SLN No. OR-010006

Rev. 1

**This label valid until December 31, 2014 or until
otherwise amended, withdrawn, cancelled or suspended.**

Active Ingredient	By Wt.
*Lactofen	23.2%
Other Ingredients	76.8%
TOTAL	100.0%
*1-(carboethoxy) ethyl 5-[2-chloro-4-(trifluoromethyl)phenoxy]-2-nitrobenzoate	
Contains 2 lb. active ingredient per gallon.	

Cobra Herbicide is not a Restricted Use Pesticide.

**KEEP OUT OF THE REACH OF CHILDREN
DANGER – PELIGRO**

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of waste. Do not apply when weather conditions favor drift from target area.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR, part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over long-sleeved shirt and long pants, chemical-resistant gloves, such as Barrier Laminate or Viton \geq 14 mils, chemical-resistant footwear plus socks, protective eyewear and chemical-resistant headgear for overhead exposure.

DIRECTIONS FOR USE

This label and the EPA registered label must be in the possession of the user at the time of mixing and pesticide application.

Follow all applicable directions, restrictions and precautions on the EPA-registered label and this label. It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM

GENERAL INFORMATION

Cobra Herbicide may be utilized as a preemergence soil-applied herbicide for control of the annual broadleaf weeds listed below immediately before transplanting or immediately following transplanting of dormant strawberry transplants. Post transplant application should be made within 48 hours of transplanting, to plants that have had soil placed over the crown of the plant to minimize direct contact of *Cobra* Herbicide with the transplants.

For preemergence control of grasses and expanded broadleaf weed control, tank mix *Cobra* Herbicide with appropriate preemergence herbicides. Examples of broadleaf weeds that *Cobra* Herbicide only partially controls include: common lambsquarters, common groundsel, shepherd's-purse and common chickweed.

Make one (1) broadcast application in the establishment year only. Do not harvest berries within one (1) year of *Cobra* Herbicide application.

Product	Product Rate	Weeds Controlled
<i>Cobra</i> Herbicide	16 to 24 fl. oz./A (0.25-0.38 lb. a.i./A)	Hairy Nightshade (<i>Solanum sarrachoides</i>) Black Nightshade (<i>Solanum nigrum</i>) Redroot Pigweed (<i>Amaranthus retroflexus</i>) Birdrape Mustard (<i>Brassica rapa</i>) Pineappleweed (<i>Matricaria matricarioides</i>)

Use Restrictions and Precautions:

1. Always read and follow all label directions when using any pesticide alone or sequentially. **The most restrictive labeling applies when using a tank-mix.**
2. Incorporate *Cobra* Herbicide with 1/4 to 1/2 inch of sprinkler-applied water following application to activate and incorporate the herbicide if rainfall is not expected within 48 hours of application.
3. *Cobra* Herbicide should not be soil incorporated with mechanical incorporation equipment.
4. Do not harvest strawberries sooner than one 1 year (365 days) after the application of *Cobra* Herbicide.

* * * * *

FIFRA Section 24(c) Registrant: VALENT U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596-8025

Cobra is a registered trademark of Valent U.S.A. Corporation