



GALIGAN® 2E HERBICIDE
EPA REG. NO. 66222-28
**USE IN FALL SEEDED NEW PLANTINGS
OF PERENNIAL RYEGRASS AND TALL
FESCUE GROWN FOR SEED
IN THE STATE OF OREGON**
OR-030034

**SUPPLEMENTAL LABELING INFORMATION FOR DISTRIBUTION AND USE
ONLY WITHIN THE STATE OF OREGON**

**This label is valid until December 31, 2012 or until otherwise amended,
withdrawn, cancelled, or suspended.**

ACTIVE INGREDIENT	% BY WT.
Oxyfluorfen: 2-chloro-1-(3-ethoxy-4-nitrophenoxy)-4-(trifluoromethyl) benzene*	22.2%
INERT INGREDIENTS:	<u>77.8%</u>
	TOTAL 100.0%

Contains petroleum distillate.
*Equivalent to 2 lbs. active ingredient per gallon.

SHAKE WELL BEFORE USING

**KEEP OUT OF REACH OF CHILDREN
WARNING-AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This label and the federal label for this product must be in the possession of the user at the time of pesticide application.

Follow all applicable directions, restrictions, and precautions on this Supplemental label and the main EPA registered label. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. This product is highly toxic to aquatic invertebrates, aquatic plants, wildlife, and fish. Use with care when applying in areas frequented by wildlife or adjacent to any body of water or wetland area. Do not apply when weather conditions favor drift or erosion from target areas.

GENERAL INFORMATION

Galigan 2E herbicide is recommended for early postemergence suppression/control of various annual broadleaf weed seedlings in fall seeded perennial ryegrass or tall fescue that has at least 1 to 2 tillers. Applications to seedling plants that have not yet tillered, may result in severe crop injury or stand loss (plant death) and is not recommended.

Note: Due to the close proximity of native prairie remnants to agricultural areas and the potential for these areas to be adversely affected by herbicides through drift or possible runoff/soil movement, it is recommended that Galigan 2E herbicide not be applied directly to native prairie habitats. It is recommended that dosage rates be maintained as the lower end of the use rate range (if effective seedling control can be achieved) in a further effort to lessen potential impacts to endangered species.

APPLICATION RATE

Apply a single application of Galigan 2E at a rate of 2 to 3 ounces (0.025 to 0.0375 lb ai) per acre, either alone or tank mixed with up to 3 pints of Nortron 4SC per acre. Some temporary crop response may occur, but is typically only a transient effect and should not adversely impact yield. **DO NOT APPLY TO NEWLY PLANTED STANDS THAT ARE UNDER STRESS FROM ANY CAUSE AS THERE IS AN ENHANCED OPPORTUNITY FOR CROP INJURY TO OCCUR.** Control from the Galigan 2E is primarily directed at emerged seedling broadleaf weeds such as speedwell and groundsel, but control or suppression of other species is possible if tank mixed with Nortron. Overlaps (2X applications) will cause significant crop injury but should not result in excessive stand losses if the crop plants are at least 1 to 2 tillers when the applications are made.

Tank mixtures of Galigan 2E with Nortron may result in enhanced crop injury. If a tank mixture is to be applied, applications should be made only to healthy, vigorous stands of perennial grasses. The decision to apply a tank mixture containing Galigan 2E herbicide is at the sole discretion of the grower and at the grower's risk.

METHOD OF APPLICATION

Apply as a broadcast application in a minimum spray volume of 20 gallons of water per acre. Use conventional ground spray equipment with flat fan spray nozzles at a minimum spray pressure of 30 psi. Do not exceed maximum spray pressure of 60 psi. Spray equipment should be calibrated carefully before each use.

MIXING DIRECTIONS

Fill the spray tank at least one-third full of clean water. With the pump and agitator running, add the recommended amount of herbicide to the spray tank. The order of addition to the spray tank should be wettable powders first, flowables second, and liquids last. Complete filling of the spray tank with water. The use of LATRON AG-98 (or comparable 80 percent active nonionic surfactant cleared for application to growing crops) may be added at a rate of 0.12 to 0.5 percent spray volume for improved control of emerged seedlings. Maintain agitation until spraying is complete.

CROP TOLERANCE

The application of Galigan 2E herbicide to fall seeded perennial ryegrass and tall fescue (that have at least 1 to 2 tillers) will result in a chlorosis (yellowing) of the vegetation within two weeks after treatment. Some crop response symptoms may be present for up to three months following application. The use of Galigan 2E herbicide may also result in a substantial reduction in vegetative growth by perennial grasses during the winter. Leaf chlorosis and reduction of vegetative growth is a typical and normal response and seed yield of healthy, vigorous perennial grasses is typically not affected by fall application of Galigan 2E herbicide. It is accepted by the grower that conditions under which seed yield may be reduced are not fully understood and that a reduction in seed yield may occur. DO NOT GRAZE FIELDS THAT HAVE BEEN TREATED WITH GALIGAN 2E HERBICIDE AS ILLEGAL RESIDUES MAY BE PRESENT IN THE VEGETATIVE FORAGE. Grazing may also magnify crop injury and reduce the seed yield. Do not apply Galigan 2E herbicide within 150 days of harvest.

WEEDS SUPPRESSED AND/OR CONTROLLED

Galigan 2E herbicide will provide control or suppression of the following weeds and volunteer crops when applied between the onset of germination and the two-leaf seedling growth stage:

COMMON NAME	SCIENTIFIC NAME
GROUNDSEL, COMMON	<i>Senecio vulgaris</i>
SPEEDWEEL	<i>Veronica Spp</i>

SPECIFIC USE RESTRICTIONS

The following use restrictions must be observed when Galigan 2E herbicide is used as recommended on this label:

- Follow the GENERAL USE RESTRICTIONS on the EPA-registered label.
- Read and observe all label directions before using. All applicable directions, restrictions, and precautions on the EPA-registered product label must be followed.
- Do not graze livestock in treated fields within 150 days of application.
- Do not apply Galigan 2E herbicide within 150 days of harvest.
- Galigan 2E herbicide should be applied only by ground application equipment.

CHEMIGATION: Do not apply this product through any type of irrigation system.

®Denotes registered trademark.

GALIGAN® is a trademark of AGAN Chemical Manufacturer, LTD., Israel.

24 (c) Registrant:

Makhteshim-Agan of North America, Inc.
4515 Falls of Neuse Rd., Suite 300
Raleigh, NC 27609

Revised: 8-9-07