



The Go To Company

P.O. Box 5569 ▲ Yuma, AZ 85366-5569 ▲ Phone (928) 783-8844 ▲ FAX (928) 343-9255

Rubigan® E.C.

EPA Reg. No. 10163-273

**24(c) Special Local Need Registration SLN No. OR-030037
For Distribution and Use only within the State of Oregon**

This label valid until December 31, 2012 or until otherwise amended, withdrawn, canceled, or suspended.

For Control of Eastern Filbert Blight in Non-bearing Hazelnuts (Filberts)

% By Wt.

ACTIVE INGREDIENT

Fenarimol: a-(2-chlorophenyl)-a-(4-chlorophenyl)-5-pyrimidinemethanol 12.0%

OTHER INGREDIENTS 88.0%

TOTAL 100.0%

Contains 1 pound active ingredient per gallon.

Contains petroleum distillate.

KEEP OUT OF REACH OF CHILDREN

WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply when weather conditions favor runoff or drift from the target area. Do not contaminate water when disposing of equipment wash waters.

DIRECTIONS FOR USE

This label and the federal label for this product must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restriction, and precautions on this Supplemental label and the main EPA-registered label. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE require for early entry to treated areas that is permitted under the Worker Protections Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves such as barrier laminate or Viton
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

GENERAL INFORMATION

Rubigan E.C. is a locally systemic fungicide that controls susceptible fungal diseases by preventing the formation of certain sterol components and other materials necessary for their growth. Rubigan E.C. is compatible with most insecticides, miticides and fungicides commonly recommended for filberts.

USE AND RATE INFORMATION

Applications of Rubigan E.C. must be made when green tissue is exposed. Apply with ground equipment in sufficient water to ensure thorough coverage. Make the first application when the first green leaf tissue is visible. Make repeat applications as necessary to maintain control. To maximize control with this product, combine with a protectant fungicide program that begins applications at bud break.

Application Rate: 4 to 8 fl. oz of Rubigan E.C. per acre.

Retreatment Interval: 7 to 14 Days.

Recommended Finished Spray Volume: 100 to 200 gallons per acre.

Minimum Spray Concentration: Do Not apply less than 4 fl oz of Rubigan E.C. per 100 gallons of spray.

Maximum Spray Concentration: Do not apply more than 8 fl oz of Rubigan E. C. per 100 gallons of spray.

RESTRICTIONS

- Do not apply to plantings that will bear harvestable nuts within 12 months of the last application or possible illegal residues may result.
- Maximum application rate per acre per application: 8 fl oz of Rubigan E.C.
- To avoid drift into bearing trees, do not apply to bearing trees interplanted in bearing orchards.
- Do not graze cover crops treated with Rubigan E.C. or harvest forage for silage or hay.
- Apply Rubigan E.C. with ground application equipment only.
- Do not apply Rubigan E.C. through any type of irrigation system.
- Observe rate restrictions.

Section 24(c) Registrant:
Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569

SLN: OR-030037 Rubigan on Hazelnuts (Filberts)