

RESTRICTED USE PESTICIDE

Due to Acute Toxicity

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

Guthion® Solupak 50% Wettable Powder Crop Insecticide

EPA Reg. No. 264-733

For use as a border treatment around alkali bee beds

SLN. No. OR-040020

24(c) Supplemental Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

UNDER A SPECIAL LOCAL NEED AUTHORIZATION UNDER FIFRA SECTION 24(c), Guthion® Solupak 50% Wettable Powder Crop Insecticide
MUST NOT BE SOLD OR DISTRIBUTED FOR USE UNDER THIS SLN AFTER DECEMBER 31, 2010.

THIS LABEL IS VALID UNTIL SEPTEMBER 30, 2012 OR UNTIL OTHERWISE AMENDED, WITHDRAWN, CANCELED OR SUSPENDED.

This label and the federal label for this product must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standards (WPS) and precautions on this supplemental label and the main EPA-registered label. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Guthion® Solupak 50%

WETTABLE POWDER CROP INSECTICIDE
IN WATER SOLUBLE PACKETS

DANGER POISON PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

ACTIVE INGREDIENT:

O,O-Dimethyl S-[(4-oxo-1,2,3-benzotriazin-3(4H)-yl)methyl]phosphorodithioate 50%

INERT INGREDIENTS:..... 50%

TOTAL 100%

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

This product is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This chemical can contaminate surface water through spray applications. Under some conditions, it may also have a high potential for runoff into surface water after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

SITE	INSECT	DOSAGE GUTHION SOLUPAK
Border treatment around alkali bee beds	Black blister beetle <i>Meloe niger</i> Blister beetle <i>Zonitis atripennis flavida</i>	See Remarks
<p>REMARKS: A ditch 4 to 6 inches wide and 6 to 10 inches deep should be dug around the alkali bee bed and then lined with plastic. Apply GUTHION Solupak at the rate of 1 lb in 8 gallons of water per 4000 feet of ditch. Initial applications should be made prior to emergence of overwintering beetles. Repeat application as necessary to maintain control. If ditch application is not feasible, apply at the rate of 20 lbs per acre in sufficient water to insure complete coverage of the area around but not on the bee bed. Repeat application at 4 to 5 day intervals during the time adult beetles are emerging.</p> <p>PITFALL TRAPS FOR SPOT TREATMENT - Install edging along bee beds and sprinkler 1-3 tablespoons (0.5 - 1.5 ounces) of Guthion Solupak 50% Wettable Powder Crop Insecticide per large pitfall trap or buried container (e.g. five-gallon plastic bucket) prior to emergence of beetles (e.g. early March). Add sufficient water into the trap at the end of season and apply spent mixture into the ground depression left after the removal of the trap.</p>		

RESTRICTIONS/PRECAUTIONS

- This product is highly toxic to bees exposed to direct application. Protective information may be obtained from your Cooperative Extension Service.
- Do not enter or allow worker entry into treated areas during the 48 hours restricted entry interval (REI) for all activities. Each 48-hour REI is increased to 72 hours in outdoor areas where average rainfall is less than 25 inches a year.
- Certain uses of Guthion® Solupak 50% Wettable Powder Crop Insecticide may be restricted by a U.S. District Court final order. You may refer to the Oregon Department of Agriculture web site at <http://egov.oregon.gov/ODA/PEST/buffers.shtml> or to the U.S. EPA web site at <http://www.epa.gov/espp/wtc/maps.htm#wtc6> or <http://www.epa.gov/espp/wtc/uselimitation.htm> for information regarding pesticides that may be impacted by the final order.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

Guthion is a registered trademark of Bayer