

# RESTRICTED USE PESTICIDE

## DUE TO ONCOGENICITY IN LABORATORY MICE\*

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



Bayer CropScience LP  
P.O. Box 12014  
2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)  
<http://www.bayercropscienceus.com>

HOELON® brand 3EC Herbicide  
For Control of Wild Oats and Annual Ryegrass (Italian)  
in Triticale.

EPA Reg. No. 264-641  
EPA SLN. No. OR-040022

## 24(c) Supplemental Label

### FOR DISTRIBUTION AND USE ONLY IN THE STATE OF OREGON

This label and the federal label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on this supplemental label and the main EPA-registered label. It is a violation of federal law to use this product in a manner inconsistent with its labeling.

THIS LABEL IS VALID UNTIL DECEMBER 31, 2012 OR UNTIL OTHERWISE AMENDED, WITHDRAWN, CANCELED OR SUSPENDED.

## HOELON® 3EC HERBICIDE

### ACTIVE INGREDIENT:

diclofop-methyl: methyl 2-[4-(2,4-dichlorophenoxy) phenoxy]propanoate..... 34.7%†

OTHER INGREDIENTS: ..... 65.3%

†Equivalent to 3.0 pounds of active ingredient per gallon.

TOTAL 100.0%

## KEEP OUT OF REACH OF CHILDREN DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

\*The Environmental Protection Agency has concluded that diclofop-methyl, the active ingredient in this product, produced tumors in laboratory mice. This effect was not observed in other species tested. The user must read and follow all precautionary statements on this label. Give particular attention to protective clothing requirements.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish. Use with care when applying in areas adjacent to any body of water. Keep out of lakes, streams, ponds, drainage basins, tidal marshes and estuaries. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor runoff or drift. Avoid direct application or drift of spray material to water surfaces. Do not apply within 100 feet of an aquatic habitat. Do not contaminate arable land and/or water when disposing of equipment washwaters.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**APPLICATION DIRECTIONS**

<b>GROUND</b>	<p>When applying HOELON® 3EC Herbicide postemergence, thorough uniform coverage of the target weeds is critical to achieving satisfactory results. The use of flat fan nozzles, spaced 10 or 20 inches apart across the boom is recommended for optimum coverage with ground equipment.</p> <p>Ground applications should be in at least 10 gallons of water carrier per acre, using a minimum of 40 psi. Ground speed for application should not exceed 10 mph.</p>
<b>AERIAL</b>	<p>Aerial applications must be applied in at least 5 gallons of water carrier per acre. Use spray nozzle tips and sufficient pressure to provide a uniform pattern and a median droplet size of 200 to 350 microns. DO NOT wide swath the application; HOELON® 3EC Herbicide is a contact material when used postemergence and highly immobile in the soil when applied preplant incorporated or preemergence. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. DO NOT make aerial applications within 100 feet of a lake, stream, drainage basin, tidal marsh, or estuary.</p> <p>Only clean water, free of suspended matter or other contaminants should be used when applying HOELON® 3EC Herbicide.</p>

**DIRECTIONS FOR THE POSTEMERGENCE USE OF HOELON® 3EC HERBICIDE ON TRITICALE**

Apply HOELON® 3EC Herbicide as a broadcast postemergence spray at the rates listed below:

Susceptible Annual Grassy Weeds	Amount of HOELON® 3EC Herbicide Per Acre (Pints) Relative to Growth Stage of Annual Grassy Weeds		
	1 – 3 Leaf	3 – 4 Leaf	5 Leaf – 2 Tillers
Annual ryegrass (Italian)..... <i>Lolium multiflorum</i>	1 1/3	1 1/3 – 2	2 – 2 2/3

**RESTRICTIONS/PRECAUTIONS**

- Make only 1 (one) application per year.
- Do not apply HOELON® 3EC less than 77 days before harvesting triticale.
- Refer to Federal label for tank mix combinations. Use only products registered for use on triticale.