

24(c) Supplemental Label
For use on the following crops grown for seed:
Broccoli, Brussels Sprouts, Cabbage, Kale, and Kohlrabi

SIMAZINE 4L
FLOWABLE HERBICIDE

EPA Reg. No. 34704-687

EPA SLN No. OR-050017

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF OREGON

This label is valid until December 31, 2012 or until otherwise amended, withdrawn, canceled, or suspended.

For use only by growers who are members of the Willamette Valley Specialty Seed Crops Association who have a signed agreement for this use filed with the Pesticide Division of the Oregon Department of Agriculture.

ACTIVE INGREDIENT:

Simazine (2-Chloro-4,6 bis (ethylamino)-s-triazine) 42.8%

INERT INGREDIENTS:..... 57.2%

TOTAL..... 100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

Simazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Simazine has been found in ground water. Users are advised not to apply simazine to sand and loamy sand soils where the water table (ground water) is close to the surface and where these soils are very permeable; i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.

Product must not be mixed or loaded within 50 feet of intermittent streams and rivers, natural or impounded lakes and reservoirs. Product must not be applied within 66 feet of points where field surface water runoff enters perennial or intermittent streams and rivers or within 200 feet of natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66 foot buffer or setback from runoff entry points must be planted to crop, or seeded with grass or other suitable crop.

Product must not be mixed or loaded, or used within 50 feet of all wells, including abandoned wells, drainage wells, and sink holes.

Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 ft. of any well are prohibited, unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rain water that may fall on the pad. Surface water shall not be allowed to either flow over or form the pad which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide to the mixing/loading sites. Additional State imposed requirements regarding well-head setbacks and operational area containment must be observed.

One of the following restrictions must be used in applying simazine to tile outletted terraced fields containing standpipes:

- Do not apply within 66 feet of standpipes in tile-outletted terraced fields.
- Apply this product to the entire tile-outletted terraced field and immediately incorporate it to a depth of 2-3 inches in the entire field.
- Apply this product to the entire tile-outletted terraced field under a no-till practice only when a high crop residue management practice is practiced. High crop residue management is described as a crop management practice where little or no crop residue is removed from the field during and after crop harvest."

DIRECTIONS FOR USE

This label and the federal label for this product must be in the possession of the user at the time of pesticide application.

Follow all other applicable directions, restrictions, and precautions on the EPA-registered label. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

APPLICATION DIRECTIONS

Apply 25.6 to 32 ounces (0.8 to 1lb. Active ingredient) per acre in the late fall when the crop is well established. Use a minimum of 20 gallons of spray solution with ground equipment. Time application to coincide with the start of the winter rainfall period to control fall-germinating annual weeds. For best results, field will have been cultivated one or more times and be relatively weed free at time of treatment. Irrigation is not required to activate the herbicide if adequate rainfall is received.

USE RESTRICTIONS/PRECAUTIONS:

- Do not apply Simazine 4L to sand, loamy sand or gravelly areas of the field. Do not apply to areas of the field where water will accumulate.
- Do not rotate to crops that are susceptible to simazine injury
- For use under this SLN label do not apply Simazine 4L through any type of irrigation system.
- Aerial application is prohibited.

Special Crop Use Restrictions

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a non-feed/non-food use by the Oregon Department of Agriculture, and is regulated by the Oregon Administrative Rule (OAR) 603-057-0535, Pesticide Use On Crops Grown For Seed. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause

the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for this crop. Consequently, no portion of this seed may be used or distributed for food or feed. This restriction pertains to, but is not limited to, green chop, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage or seed screenings, and to the grazing of the crop field, stubble or re-growth. All seed screenings shall be disposed of in such a manner that screenings cannot be distributed or used for food or feed purposes, as indicated in OAR 603-057-0535. Additional regulations concerning seed screenings are stated in OAR 603-057-0535.

Any seed from the field treated with this pesticide product shall bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

“This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. This seed, in whole, as sprouts, or in any form may not be used for human consumption or animal feed. Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration, the Oregon Department of Agriculture and other regulatory agencies.”