

RESTRICTED USE PESTICIDE

Due to acute oral, dermal, inhalation, eye, and skin toxicity

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

Di-Syston® 8 Emulsifiable Systemic Insecticide
For Use on Commercially Grown Easter Lilies

EPA Reg. No. 264-734
EPA SLN. No. OR-050024

24(c) Supplemental Label

FOR DISTRIBUTION AND USE ONLY WITHIN CURRY COUNTY IN THE STATE OF OREGON

This label and the federal label for this product must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on this supplemental label and the main EPA-registered label. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS LABEL IS VALID UNTIL DECEMBER 31, 2012 OR UNTIL OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

Di-Syston® 8 Emulsifiable Systemic Insecticide

ACTIVE INGREDIENT:

Disulfoton, O,O-Diethyl S-[2-(ethylthio)ethyl]-phosphorodithioate..... 85%

OTHER INGREDIENTS: 15%

Contains petroleum distillates 100%



**STOP – Read the label before use.
KEEP OUT OF REACH OF CHILDREN
DANGER POISON
PELIGRO**



Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to birds, mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to fish in neighboring areas. Do not contaminate water when cleaning equipment or when disposing of equipment washwaters. Cover or incorporate spills.

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months. Poor draining soils and soils with shallow water tables are prone to produce runoff that contains this product. Disulfoton is known to leach through the soils into ground water under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

A level, well-maintained 25 foot vegetative buffer strip between areas to which this product is applied and permanent surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Refer to the

March 2000 USDA Natural Resources Conservation Service document: *Conservation Buffers to Reduce Pesticide Losses* for additional guidance. Runoff of this product will be reduced by avoiding application when rainfall is forecasted to occur within 48 hours.

This pesticide is extremely toxic to birds, mammal, fish and aquatic invertebrates. Di-Syston 8 Emulsifiable Systemic Insecticide should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may contact the Oregon Department of Fish & Wildlife, National marine Fisheries Services, or US Fish & Wildlife Service for information on listed threatened or endangered species. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift.

Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

RECOMMENDED APPLICATIONS			
CROP	INSECT	DOSAGE	REMARKS
Easter Lilies	Aphids Lace bug Leafhoppers Mites Thrips Whiteflies	0.53 to 1.1 fl oz per 100 ft of row	Apply in the furrow over the bulbs after they are set, but before covering with soil. The higher dosage is recommended for heavy organic or muck soils.

RESTRICTIONS/PRECAUTIONS

- Maximum number of applications: 1 per year
- Do not apply this product through any type of irrigation system.
- Do not enter or allow worker entry into treated areas during the restricted entry interval of 48 hours. The REI is 72 hours in areas where average rainfall is less than 25 inches a year.
- Do not apply within a minimum of 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds. A level, well-maintained 25 foot vegetative buffer strip between areas to which this product is applied and permanent surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Refer to the March 2000 USDA Natural Resources Conservation Service document: *Conservation Buffers to Reduce Pesticide Losses* for additional guidance. Runoff of this product will be reduced by avoiding application when rainfall is forecasted to occur within 48 hours.

Certain uses of disulfoton may be restricted by a U.S. District Court final order. You may refer to the Oregon Department of Agriculture web site at <http://egov.oregon.gov/ODA/PEST/buffers.shtml> or to <http://www.epa.gov/espp/litstatus/wtc/maps.htm> for information regarding pesticides that may be impacted by the final order.

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

Di-Syston is a registered trademark of Bayer