



Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

LSP Flowable Fungicide
 For Use on Crimson Clover grown for Seed

 EPA Reg. No. 264-948
 EPA SLN. No. OR-060003

24(c) Supplemental Label

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF OREGON

This label and the federal label for this product must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on this supplemental label and the main EPA-registered label. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS LABEL IS VALID UNTIL DECEMBER 31, 2013 OR UNTIL OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

LSP Flowable Fungicide

ACTIVE INGREDIENT:

Thiabendazole: 2-(4-thiazolyl) benzimidazole30%

INERT INGREDIENTS:70%

TOTAL 100%

Contains Petroleum Distillate

Note to Physician: Contains petroleum distillate – vomiting may cause aspiration pneumonia.

In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577.
 Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

KEEP OUT OF REACH OF CHILDREN CAUTION

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on agricultural establishments in hopper-box, planter-box, slurry-box, or other seed treatment applications at or immediately before planting.

GENERAL: LSP is a systemic fungicide especially formulated as a liquid suspension seed protectant for use as a spray mist or slurry seed treatment.

SEED TREATMENT: Clean seed treatment equipment thoroughly before using LSP. This product has no vapor action; therefore, it is essential that thorough coverage of the seed be obtained. In this regard, LSP is recommended to be dispersed with water to aid distribution on the seed. When dispersed with water, however, the resulting suspension must be kept under constant agitation.

DIRECTIONS

CRIMSON CLOVER (GROWN FOR SEED)

CROP/SITE	TARGET PEST:	RATE and METHOD OF APPLICATION
Crimson Clover (Grown for Seed)	For seed use in control of seedborne Northern Anthracnose (<i>Kabatiella caulivora</i>)	For ready-mix or slurry seed treaters, mix 13.0 fl.oz. of LSP Flowable Fungicide in sufficient water to provide adequate coverage and apply to 100 lbs. of seed. Keep dispersion under constant agitation. Allow seed to dry thoroughly before planting. Rhizobium spp. for nitrogen fixation should be applied immediately prior to planting (within 4 hours). If inoculant is left in contact with treated seed for long periods of time, re-inoculation is required.

Restrictions/Precautions:

USE PRECAUTION: When using formulations that do not contain dye, to comply with 21 CFR Chapter 1, Section 2.25, all grain seed treated with an economic poison must be colored with an approved dye (40 CFR 180.1001) to distinguish and prevent subsequent and inadvertent use as a food for man or feed for animals.

COMPATIBILITY: LSP should not be tank mixed with any pesticide unless adequate physical compatibility and phytotoxicity tests have been completed and proved to be satisfactory.

SEED LABELING: The directly treated seed must be labeled, "This seed is treated with Thiabendazole at the manufacturer's recommended rate. Not for human or animal consumption. Do not use treated seed for food, feed, or oil purposes." Add appropriate information for any other chemical treatments.

Special Crop Use Restrictions

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a nonfeed/non-food use by the Oregon Department of Agriculture, and is regulated by Oregon Administrative Rule (OAR) 603-057-0535, Pesticide Use On Crops Grown For Seed. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for this crop. Consequently, no portion of this seed crop may be used or distributed for food or feed for 1 year (365 days) after the last application of this product. This restriction pertains to, but is not limited to: green chop, forage, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage or seed screenings, and to the grazing of the crop field, stubble or regrowth. All seed screenings shall be disposed of in such a manner that the screenings cannot be distributed or used for food or feed purposes, as indicated in OAR 603-057-0535. Additional regulations concerning seed screenings are stated in OAR 603-057-0535.

Any seed from a field treated with this pesticide product shall bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

"This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. This seed, in whole, as sprouts, or in any form, may not be used for human consumption or animal feed. Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration, the Oregon Department of Agriculture and other regulatory agencies."

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).