



Supplemental Labeling

SPECIAL LOCAL NEED FOR DISTRIBUTION AND USE ONLY WITHIN OREGON

EPA Reg. No. 241-418

EPA SLN No.: OR-060007

FOR WEED CONTROL IN FIRST YEAR NON-BEARING STRAWBERRIES

ACTIVE INGREDIENT:

pendimethalin (N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine).....38.7%

INERT INGREDIENTS:.....61.3%

TOTAL.....100.0%

This label is valid until December 31, 2010 or until otherwise amended, withdrawn, canceled or suspended.

NOTICE

BASF CORPORATION ("BASF") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PROWL H₂O HERBICIDE AS AN AID IN CONTROLLING WEEDS IN FIRST YEAR NON-BEARING STRAWBERRIES. BASF DOES NOT WARRANT THE USE OF PROWL ON FIRST YEAR NON-BEARING STRAWBERRIES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered main product label for Prowl® H₂O herbicide, EPA Reg. No. 241-418.

This labeling must be in the possession of the user at the time of application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Observe all cautions and limitations on this label and on the labels of products used in combination with **Prowl H₂O**. Do not use **Prowl H₂O** other than in accordance with the instructions set forth on this label. The use of **Prowl H₂O** not consistent with this label may result in injury to crops. Keep containers closed to avoid spills and contamination.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.



GENERAL INFORMATION

Prowl® H₂O herbicide may be applied for pre-emergence weed control of most annual grasses and certain broadleaf weeds as they germinate, but it will not control established weeds. Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend it over a long period of time may reduce weed control.

Over-application can result in crop stand loss, crop injury, or rotational crop injury.

Uneven application or improper soil incorporation can decrease weed control or cause crop injury. Soil incorporation deeper than recommended can reduce weed control.

Use of **Prowl H₂O** in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

Refer to the **Prowl H₂O** main product label for a complete listing of weeds controlled.

APPLICATION INSTRUCTIONS

Stunting, reduced growth, or reduction in daughter plants may occur with this use. Uniformly broadcast apply **Prowl H₂O** preemergence prior to transplanting strawberries. Make one application in 10 or more gallons of water per acre with ground equipment.

APPLICATION RATES

**Broadcast Rate Per Acre of Prowl H₂O
Used In First Year Non-Bearing Strawberries
(Pints Per Acre)**

Soil Texture	<3% Organic Matter	>3% Organic Matter
Coarse	1.5 – 2.0 pts.	2.0 – 3.0 pts.
Medium	2.0 – 2.5 pts.	2.0 – 3.0 pts.
Fine	2.0 – 3.0 pts.	2.5 – 3.5 pts.

RESTRICTIONS AND LIMITATIONS

For use under this SLN label **DO NOT** apply this product through any type of irrigation system, or by air.

DO NOT use on peat or muck soils.

Prowl H₂O is most effective when incorporated in the weed germination zone by rainfall or irrigation.

DO NOT exceed the maximum labeled rate for a given soil texture.

DO NOT plant winter wheat or winter barley in the fall following **Prowl H₂O** application at rates above 3.0 pints per acre.

NOTE: If loss of strawberry stand occurs due to adverse weather conditions, any crop registered for **Prowl H₂O** preplant incorporated use can be replanted the same year into **Prowl H₂O** treated soil without adverse effects. If replanting is necessary, soil should not be reworked deeper than the **Prowl H₂O** treated zone.

DO NOT harvest for food or feed any portion of the strawberry plant within 1 year (365 days) of **Prowl H₂O** application.

©2005 BASF Corporation
All rights reserved

Prowl is a registered trademark of BASF.

0000241-00418.20051018c.NVA 2005-04-195-0288

24(c) Registrant:

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709


The Chemical Company