

RESTRICTED USE PESTICIDE

ACUTE DERMAL TOXICITY

For retail sale to and use only by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's certification.



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

MOCAP® EC Nematicide-Insecticide
For Use on Snap Beans

EPA Reg. No. 264-458

EPA SLN. No. OR-060010

24(c) Supplemental Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

This label and the federal label for this product must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on this supplemental label and the main EPA-registered label. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS LABEL IS VALID UNTIL DECEMBER 31, 2009 OR UNTIL OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

MOCAP® EC Nematicide-Insecticide

DANGER POISON

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

DANGER

Poisonous if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. Do not get into eyes, on skin, or on clothing. Do not breathe fumes or spray mist. Do not contaminate feed or food.

SYMPTOMS OF POISONING: Nausea, vomiting, abdominal cramps, diarrhea, excessive salivation, headache, dizziness, weakness, blurring or dimness of vision, excessive tearing, loss of muscular coordination, slurring of speech, twitching of muscles (especially of tongue and eyelids), mental confusion, disorientation, drowsiness, difficulty in breathing (chest tightness), runny nose.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms (fish and invertebrates) and extremely toxic to birds. **Cover or disc spill areas.** Birds in treated areas may be killed. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment wash waters.

Do not apply within 140 feet of inland freshwater habitats.

Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

Drift and runoff may be hazardous to aquatic organisms in neighboring areas.

DIRECTIONS FOR USE
Beans (snap)

RATE OF MOCAP® EC

APPLICATION TIMING	PESTS CONTROLLED	BANDED FL.OZ./ 1000 ROW FEET	BROADCAST GALLONS PER ACRE	APPLICATION DIRECTIONS
at planting or up to 3 days before planting	nematodes garden symphylans	2 to 3.9	1 to 1 1/3	Band: Apply by shanking or injecting into the soil about 1 to 2 inches to the side of the seed. Do not use as a seed furrow treatment or allow spray to contact the seed or phytotoxicity may occur. Broadcast: Apply using the indicated broadcast rate per acre and immediately mix with the top 2 to 4 inches of soil.
	garden symphylans	2 to 3.9	1/3 to 1/2 (or 1 1/3 to 2 quarts)	
Note: Beans grown in treated soil may remain green longer than other beans and mature 1 to 2 weeks later. Make only one application per crop.				

RESTRICTIONS AND PRECAUTIONS

- Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks, and recreation areas, non-target crops, aquatic and wetland areas, woodlands, pastures, rangelands or animals.
- Apply through ground equipment only; Do not apply by air or through irrigation systems.
- Mixer/loader/applicators must wear Personal Protective Equipment (PPE), including respirators. Mixers/loaders/applicators must be observed for any sign of central nervous system disorders (i.e. nausea, diarrhea, vomiting, sweating, salivation, and lacrimation). Should these symptoms occur, workers should immediately be seen by a physician and have their cholinesterase levels monitored.
- Do not graze livestock in treated areas.
- Post "Danger, Skull and Crossbones" warning signs at all entrances to treated areas.
- To protect endangered avian species, the product should not be used in areas of known species occurrence and during the breeding season in those areas.

Certain uses of ethoprop may be restricted by a U.S. District Court final order. You may refer to the Oregon Department of Agriculture web site at <http://oregon.gov/ODA/PEST/buffers.shtml> or to the U.S. EPA web site at <http://www.epa.gov/espp/wtc/maps.htm#wtc6> for information regarding pesticides that may be impacted by the final order.

**For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)**

or

Visit our worldwide web site at www.bayercropscienceus.com

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*