

# RESTRICTED USE PESTICIDE

DUE TO ACUTE ORAL, DERMAL AND INHALATION TOXICITY

For retail sale to and use only by Certified Applicators or persons under the direct supervision of a Certified Applicator, and only for those uses covered by the Certified Applicator's certification.



Bayer CropScience

Bayer CropScience LP  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)  
<http://www.bayercropscienceus.com>

MOCAP® EC Nematicide-Insecticide

For Use on Sugar Beets grown for seed.

EPA Reg. No. 264-458

EPA SLN. No. OR-o60024

**24(c) Supplemental Label**

## FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

### (West of the Cascade Mountain Range)

This label and the federal label for this product must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on this supplemental label and the main EPA-registered label. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS LABEL IS VALID UNTIL DECEMBER 31, 2014 OR UNTIL OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

## MOCAP® EC Nematicide-Insecticide

DANGER POISON

### PRECAUTIONARY STATEMENTS

#### HAZARD TO HUMANS AND DOMESTIC ANIMALS

##### DANGER

Poisonous if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. Do not get into eyes, on skin, or on clothing. Do not breathe fumes or spray mist. Do not contaminate feed or food. Corrosive. Causes irreversible eye damage.

**SYMPTOMS OF POISONING:** Nausea, vomiting, abdominal cramps, diarrhea, excessive salivation, headache, dizziness, weakness, blurring or dimness of vision, excessive tearing, loss of muscular coordination, slurring of speech, twitching of muscles (especially of tongue and eyelids), mental confusion, disorientation, drowsiness, difficulty in breathing (chest tightness), runny nose.

### ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms (fish and invertebrates) and extremely toxic to birds. **Cover or disc spill areas.** Birds in treated areas may be killed. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment wash waters.

Do not apply within 140 feet of inland freshwater habitats.

Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

Drift and runoff may be hazardous to aquatic organisms in neighboring areas.

## Sugar Beets grown for seed

APPLICATION TIMING	PEST CONTROLLED	RATE OF MOCAP® EC BROADCAST QUARTS PER ACRE	APPLICATION DIRECTIONS
Prior to seeding or transplanting	garden symphylans	1 1/3 to 2	<b>Broadcast &amp; Incorporate</b> Mix product with sufficient water and broadcast apply immediately ahead of equipment such as a double disc or rotary cultivator to thoroughly incorporate MOCAP EC solution into the top 2 to 4 inches of soil.
After Seeding Emergence or After transplanting seedlings. Suppression only	garden symphylans	1 1/3 to 2	<b>Shanking or Injecting</b> Shank or inject solution 1 to 2 inches from plants and to a depth of 2 to 4 inches on both sides of seedlings or transplants.  Mix with sufficient water and inject product to side of plants close to but not in contact with the roots. Follow with irrigation.  Do not apply to a seed furrow or allow spray to contact plants. Do not shank/ inject into excessively wet or dry soils.
<b>Note:</b> Sufficient soil moisture should be present at application for symphylans to be active in the top 4 to 6 inches of the soil profile. Volatilization may happen in very dry soils. Do not apply to dry soils.			

### RESTRICTIONS AND PRECAUTIONS

- Do not make more than one application to the same field per crop season.
- Do not apply by any method of application not stated on this label. The specific application directions must be followed for each application timing.
- This product must be mechanically soil incorporated, shanked or injected. It is not sufficient to simply spray MOCAP EC on the soil surface and irrigate.
- Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks, and recreation areas, non-target crops, aquatic and wetland areas, woodlands, pastures, rangelands or animals.
- Apply only through motorized ground equipment. Do not apply by air or through irrigation systems.
- Mixer/loader/applicators must wear Personal Protective Equipment (PPE) specified on the main EPA-registered label. In addition, mixers and loaders must use a mechanical transfer system as described on the main EPA-registered label. Applicators using motorized ground equipment must use an enclosed cab as described on the main EPA-registered label.
- Do not graze livestock in treated areas.
- Post "Danger, Skull and Crossbones" warning signs at all entrances to treated areas.  
To protect endangered avian species, the product should not be used in areas of known species occurrence and during the breeding season in those areas.

You are advised that the use of this product in certain portions of California, Oregon and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et. Al. v. EPA, CO1-0132C, (E.D.WA). For further information please refer to the following EPA website: <http://www.epa.gov.espp/wtc/uselimitation.htm>.

You may also refer to the Oregon Department of Agriculture web site at: <http://egov.oregon.gov/ODA/PEST/buffers.shtml>.

### **Special Crop Use Restrictions**

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a nonfeed/non-food use by the Oregon Department of Agriculture, and is regulated by Oregon Administrative Rule (OAR) 603-057-0535, Pesticide Use On Crops Grown For Seed. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for this crop. Consequently, no portion of this seed crop may be used or distributed for food or feed for 1 year (365 days) after the last application of this product. This restriction pertains to, but is not limited to: green chop, forage, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage or seed screenings, and to the grazing of the crop field, stubble or regrowth. All seed screenings shall be disposed of in such a manner that the screenings cannot be distributed or used for food or feed purposes, as indicated in OAR 603-057- 0535. Additional regulations concerning seed screenings are stated in OAR 603-057-0535. Any seed from a field treated with this pesticide product shall bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

"This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. This seed, in whole, as sprouts, or in any form, may not be used for human consumption or animal feed. Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration, the Oregon Department of Agriculture and other regulatory agencies."

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577  
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at [www.bayercropscienceus.com](http://www.bayercropscienceus.com)

*As with any crop-protection product, always read and follow label directions.  
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*