



**For the Control of  
Spider Mites in Potatoes**

**FIFRA Section 24(c) Supplemental Label  
ACRAMITE®-4SC Agricultural Miticide  
FOR DISTRIBUTION AND USE  
ONLY WITHIN THE STATE OF OREGON**

This label for Acramite-4SC expires and is valid until December 31, 2012 or until otherwise amended, withdrawn, canceled, or suspended.

**EPA Reg. No. 400-514 EPA SLN No. OR-070019**

Active Ingredient: (% by weight)  
bifenazate: hydrazine carboxylic acid, 2-(4-methoxy-  
[1,1'-biphenyl]-3-yl) 1-methylethyl ester ..... 43.2%  
Inert Ingredients: ..... 56.8%  
Total ..... 100.0%  
Contains 4 lbs. active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this supplemental label and the main EPA-registered label.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to birds, estuarine/marine invertebrates, and fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treatment area.

**Purpose of this Supplemental Label: Add Chemigation Instructions**

**Plant-Back Restrictions:** Do not plant another crop within 30 days after last ACRAMITE application due to chances of bifenazate residue showing up in rotational crops. Exception: If bifenazate is registered for use on a crop, there are no plant-back restrictions.

**DIRECTIONS FOR USE:  
POTATOES**

Target Pests	Application Rates Fl. Oz. Per Acre	Minimum Gallons Per Acre Chemigation	Application Timing
Two spotted Spider Mite	16-24	Apply in 0.1 to 0.2 acre inches of water	To provide maximum residual control, application should be made as soon as mites appear. Use the lower rate where mite infestations are light. The higher rate may be required for heavy infestations or for extended residual control.

**DIRECTIONS FOR USE** table exhibits application rate range. Only one application may be made per year. Sprinkler systems should be operated at 80 to 100% during treatment application to apply the minimum amount of water possible. However, Acramite-4SC is not systemic in action; therefore complete coverage of both upper and lower leaf surfaces is necessary for effective control.

### **CHEMIGATION USE PRECAUTIONS FOR POTATOES**

- A. Apply this product only through sprinkler systems, including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system.
- B. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.
- C. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- D. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.
- E. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make the necessary adjustments should the need arise.
- F. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- G. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- H. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- I. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- J. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- K. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- L. Do not apply when wind speed favors drift beyond the area intended for treatment.
- M. Constant agitation must be maintained in the chemical supply tank during the entire period of miticide application.
- N. Inject the product with a positive replacement pump into the main line ahead of a right angle turn, to insure adequate mixing.
- O. Application of more than label recommended quantities of irrigation water per acre may result in decreased product performance by removing the chemical from the zone of effectiveness.
- P. Do not apply when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.
- Q. Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of more dilute mixture per hour. Pesticide should be applied continuously for the duration of the water addition.
- R. Where sprinkler irrigation patterns do not overlap sufficiently unacceptable mite control may result. Where sprinkler distribution patterns overlap excessively crop injury may result.

**Special Crop Use Restrictions:**

This pesticide is toxic to birds, estuarine/marine invertebrates, and fish. Acramite-4SC should not be used under this SLN label where impact on listed threatened or endangered species is likely.

**SLN 24c – Registrant:**

**Chemtura USA Corporation**

**Middlebury, CT 06749**

Acramite-4SC is a registered trademark of Chemtura Corporation

Rev. 06/29/2007