



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

Axiom® DF Herbicide
EPA Reg. No. 264-766
SLN No. OR-080027

For use in annual ryegrass grasses
grown for seed fields.

24(c) Supplemental Label

**FOR DISTRIBUTION AND USE ONLY WEST OF THE CASCADE MOUNTAIN RANGE
IN THE STATE OF OREGON**

This label and the federal label for this product must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on this supplemental label and the main EPA-registered label. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS LABEL IS VALID UNTIL DECEMBER 31, 2013 OR UNTIL OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

Axiom® DF Herbicide

ACTIVE INGREDIENTS:

Flufenacet

N-(4-Fluorophenyl)-*N*-(1-methylethyl)-2-[[5-(trifluoromethyl)-1,3,4-thiadiazol-2-yl]-oxy]acetamide..... 54.4%

Metribuzin:

4-Amino-6-(1,1-dimethylethyl)-3-(methylthio)-1,2,4-triazin-5(4*H*)-one..... 13.6%

INERT INGREDIENTS: 32.0%

100.0%

**STOP - Read the label before use
KEEP OUT OF REACH OF CHILDREN
CAUTION**

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters.

Do not allow sprays to drift onto adjacent desirable plants.

Metribuzin has properties that may result in ground water contamination. Application in areas where soils are permeable or coarse and groundwater is near the surface could result in groundwater contamination.

Ground Water: Flufenacet is known to leach through soil into ground water under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water: Flufenacet and metribuzin residues can contaminate surface water through spray drift. Under some conditions, they also have high potential for runoff into surface water (primarily via dissolution in runoff water or erosion), for several to many months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlying tile drainage systems that drain to surface waters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass

Bromus*

Italian (annual) ryegrass

Rattail Fescue

Rough Bluegrass

Volunteers of the established crop

* indicates suppression.

RECOMMENDED APPLICATIONS

Row Spray Application - For Use in Annual Ryegrass grown for seed.

Crop	Axiom® DF Rate (oz. per acre)
Annual Ryegrass	8 to 10
REMARKS Apply specified rate per treated acre as a band spray application to remove unwanted excessive volunteer annual ryegrass, and weeds such as annual bluegrass from the treated bands. Apply using shielded sprayers to prevent drift to unsprayed rows where the desired seed grass crop is grown. Use with a registered contact herbicide such as glyphosate to remove volunteers and weeds which have germinated and emerged prior to treatment.	

SPECIAL PRECAUTIONS:

Do not apply this product through any type of irrigation equipment. Do not apply in tank-mixture with postemergence herbicides, such as Diuron. Application of oxyfluorfen or photosynthetic inhibiting postemergence herbicides within 4 weeks of Axiom® applications may cause crop damage.

Weed control may be reduced if excessive straw from the previous crop is present at Axiom® application or if insufficient rainfall/irrigation occurs after application.

RESTRICTIONS:

- Do not allow animals to graze treated fields for a minimum of 30 days following application.
- Do not apply by aerial application.
- Do not make more than one application per season.
- Do not apply more than 10 oz of Axiom DF per acre per use season.
- For all metribuzin products used, do not apply more than a total of 1 lb ai/A per use season of metribuzin.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*