



FIFRA 24(c) REGISTRATION

**REGISTRATION FOR SPECIAL LOCAL NEED
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON**

**EVITO 480 SC Fungicide
EPA REG. NO. 66330-64 EPA SLN No: OR-080034**

This label valid until December 31, 2013 or until otherwise amended, withdrawn, cancelled, or suspended.

FOR USE ON POTATO (BANDED AND IN-FURROW APPLICATIONS)

ACTIVE INGREDIENT: Fluoxastrobin: [(1 <i>E</i>)-[2-[[6-(2-Chlorophenoxy)-5-fluoro-4-pyrimidinyl]oxy]phenyl] 5,6-dihydro-1,4,2-dioxazin-3-yl) methanone-O-methyloxime]	40.3%
INERT INGREDIENTS:	59.7%
TOTAL:	100.0%

DIRECTIONS FOR USE

- This label and the federal label for this product must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, and precautions on this Supplemental label and the main EPA registered label.
- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. The active ingredient in this product can be persistent for several months or longer. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark, or other sensitive areas that may be exposed to spray drift. Do not contaminate water when disposing of equipment washwater or rinsate.

POTATO

Soilborne / Seedling Diseases Controlled	Rate to Use	Application Timing
Black scurf (<i>Rhizoctonia solani</i>)	0.16 – 0.24 fl oz./1000 row feet	See directions and rates under SOILBORNE / SEEDLING DISEASE CONTROL.

SOILBORNE / SEEDLING DISEASE CONTROL

Evito can provide control of soilborne diseases, including Black scurf, if applied early in the growing season. Specific applications for soilborne diseases include in-furrow applications or banded applications applied over the row, either shortly after plant emergence or during herbicide applications or cultivation. These applications will provide control of pre- or post-emergence damping off and diseases that infect plants at the soil-plant interface. The use of either type of application depends on the cultural practices in the region. In some locations, one type of application may provide better disease control than the other, depending on the timing of the disease epidemic. Seedling diseases are generally controlled by in-furrow applications while banded applications are more effective against soilborne diseases that develop later in the season. Consult your local expert for guidance regarding application type.

Banded applications: Apply Evito prior to infection as a directed spray to the soil, using single or multiple nozzles, adjusted to provide thorough coverage of the lower stems and the soil surface surrounding the plants. Band width should be limited to 7 inches or less. Apply Evito at a rate of 0.16-0.24 fl. oz. product/1000 row feet. These applications come into contact with the foliage and are counted as foliar applications when considering resistance management. They may be applied during cultivation or hilling operations to provide soil incorporation.

In-furrow applications: Apply Evito as an in-furrow spray in 3-7 gallons of water at planting. Mount the spray nozzle so the spray is directed into the furrow just before the seed are covered. Use the higher rate when the weather conditions are expected to be conducive for disease development, if the field has a history of *Pythium* problems, or if minimum/low till programs are in place.

IN-FURROW APPLICATION RATES

RATE PER 1000 ROW FEET	PRODUCT PER ACRE (fl. oz.)						
	22" rows	30" rows	32" rows	34" rows	36" rows	38" rows	40" rows
fl. oz. product							
0.16	3.8	2.8	2.6	2.5	2.3	2.2	2.1
0.24	-	-	3.9	3.7	3.5	3.3	3.1

40" = 13,068 row ft., 38" = 13,754 row ft., 36" = 14,520 row ft., 34" = 15,374 row ft., 32" = 16,315 row ft., 30" = 17,424 row ft., and 22" = 23,760 row ft./Acre

RESTRICTIONS / PRECAUTIONS:

- Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.
- Do not apply more than 22.8 fl oz (0.72 lbs a.i.) of EVITO 480 SC per acre per year (including banded and in-furrow applications).
- There is a maximum number of 6 applications per season, and a minimum interval of 7 days between applications.
- Do not use the 0.24 fl. oz/1000 row feet rate on 22 or 30 inch rows.
- Do not apply EVITO 480 SC within 7 days of harvest.
- For use under this SLN label do not apply this product through any type of irrigation system.
- This pesticide is toxic to fish and aquatic invertebrates.

CONDITIONS OF SALE

Arysta LifeScience North America Corporation ("Arysta") warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use.

This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Arysta.

ARYSTA DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE OR MERCHANTABILITY. TO THE EXTENT PERMITTED BY LAW, SELLER SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM OR IN CONNECTION WITH THE MANUFACTURE, SALE, DELIVERY, USE, HANDLING OR STORAGE OF THIS PRODUCT. TO THE EXTENT PERMITTED BY LAW, SELLER'S LIABILITY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. ARYSTA DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.

Critical and unforeseeable factors beyond Arysta's control prevent Arysta from eliminating all risks in connection with the use of this product. Such risks include, but are not limited to, damage to plants and crops to which the product is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated on the label and even though label directions are followed. Except as stated above, to the extent permitted by law, by purchasing, accepting and using this product, the Buyer and User acknowledge and assume all risks and liability resulting from handling, storage, and use of this product.

EVITO is a trademark of Arysta LifeScience North America Corporation

24(c) Registrant:

**Arysta LifeScience North America Corporation
15401 Weston Parkway, Suite 150
Cary, NC 27513**

Rev. 09-23-2008