

FIFRA SECTION 24(c) SUPPLEMENTAL LABELING
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON



Vitaflo[®] 280

FLOWABLE FUNGICIDE

For Use on Grass Seed - For Export Only

EPA REG. NO. 400-564

EPA SLN NO. OR-080036

This label and the federal label for this product must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on this supplemental label and the main EPA-registered label. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

THIS LABEL IS VALID UNTIL December 31, 2013 OR UNTIL OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

Active Ingredients:	
Carboxin (5,6-dihydro-2-methyl-M-phenyl-1,4-oxathiin-3-carboxamide)	14.9%*
Thiram (Tetramethylthiuram disulfide).....	13.2%*
Other Ingredients:	71.9%
Total:.....	100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

- | | |
|--------------------------------|---|
| IF IN EYES: | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lens, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice. |
| IF SWALLOWED: | <ul style="list-style-type: none">• Immediately call a poison control center or doctor.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give any liquid to the person.• Do not give anything by mouth to an unconscious person. |
| IF INHALED: | <ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible.• Call a poison control center or doctor for further treatment advice. |
| IF ON SKIN OR CLOTHING: | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice. |

NOTE TO PHYSICIAN: Contains petroleum distillate - vomiting may cause aspiration pneumonia.

EMERGENCY ASSISTANCE: Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

EMERGENCY PHONE	800-292-5898
SAFETY DATA AND INFORMATION	866-430-2775
TRANSPORTATION EMERGENCY (CHEMTREC)	800-424-9300

DIRECTIONS FOR USE

This label and the federal label for this product must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on this label and the main EPA-registered label. It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

GENERAL INFORMATION

VITAFLO™ 280 aids in the control of seed decay, damping-off and seedling blight.

Use only as a grass seed treatment for grass seed being exported to countries requiring treatment of carboxin before importation. Users should verify actual registered rate in country of destination and/or use. This label is based solely on treatment requirements made by countries requiring carboxin to be applied to imported grass seed. Chemtura Corporation makes no claims as to the effect of this product on disease control or seed germination. All such risks should be assumed by the Buyer.

APPLICATION DIRECTIONS – COMMERCIAL SEED TREATMENT

VITAFLO 280 is a seed treatment which should be applied as a slurry treatment by machine.

Application should be made using mechanical slurry or mist-type treatment equipment. Use 3.3 to 12 fluid ounces VITAFLO 280 per 100 pounds of seed.

002

24(c) Registrant:
Chemtura Corporation
199 Benson Road
Middlebury, Connecticut 06749

SLN No. OR-080036

In most cases, VITAFLO 280 is ready to use and may be applied to seed as is; however, dilution with water may be appropriate for some types of treaters and/or when treating under dry and/or hot conditions, to achieve more uniform product to seed coverage. VITAFLO 280 and water should be mixed thoroughly into a slurry prior to treating seed. Recalibrate treating equipment to compensate for the addition of water. The amount of dilution water that is most suitable depends on the seed, treater type and temperature. Contact your local Chemtura representative or supplier for specific recommendations.

Seed should be sound and well cured prior to treatment. Care must be exercised in the handling of treated seed.

USE RESTRICTIONS:

- Do not use treated seed for food or feed purposes.
- Treated seed cannot be planted in the United States and must be exported.

COMPATIBILITY: VITAFLO 280 can be used in combination with other flowable seed protectant fungicides and insecticides. Consult a seed treatment specialist regarding combining of products with VITAFLO 280 for specific seed crop allocations and slurry compatibility. Tank mixes should be pre-tested to determine physical compatibility between formulations, and to assure there will be no adverse phytotoxic reactions.

SEED LABELING (Commercial Seed): Treated seed must be labeled, For Export Only – Not to be sold or offered for sale in the United States. The U.S. Seed Act requires that treated seed be labeled with the following information: This seed treated with Carboxin and Thiram. Do not use treated seed for food, feed, or oil purposes.

NOTE: Treated seeds exposed on soil surface may be hazardous to birds. Cover or collect spilled seeds.