

SUPPLEMENTAL LABELING



Simazine 4L

Herbicide

(EPA REG. NO. 19713-60)

(EPA SLN NO. OR-090014)

**FOR USE AS HERBICIDE ON THE FOLLOWING CROPS GROWN FOR SEED:
BROCCOLI, BRUSSELS SPROUT, CABBAGE, CHINESE MUSTARD, FALL PLANTED
RADISH, KALE, KOHLRABI, AND TURNIP**

For Distribution and Use Only in the State of Oregon

INTENDED FOR USE ONLY BY THE SPECIALTY SEED GROWERS OF WESTERN OREGON (SSGWO) AND MEMBERS OF THE WILLAMETTE SPECIALTY SEED CROPS ASSOCIATION

ACTIVE INGREDIENT:

Simazine: 42.1%

OTHER INGREDIENTS: 57.9%

TOTAL: 100.0%

This product contains 4 pounds of Simazine per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

ENVIRONMENTAL HAZARDS

Simazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Simazine has been found in ground water. Users are advised not to apply simazine to sand and loamy sand soils where the water table (ground water) is close to the surface and where these soils are very permeable; i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This pesticide is toxic to aquatic invertebrates. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

Product may not be mixed/loaded, or used within 50 feet of all wells, including abandoned wells, drainage wells, and sink holes. Consult the federal label for additional restrictions on mixing, loading, rinsing or washing of this product.

Product must not be mixed or loaded within 50 feet of intermittent streams or rivers, natural or impounded lakes and reservoirs. Product must not be applied within 66 feet of the points where field surface water runoff enters perennial or intermittent streams and rivers or within 200 feet of natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66-foot buffer or setback from runoff points must be planted to crop or seeded with grass or other suitable crop.



Manufactured By:
Drexel Chemical Company
P.O. BOX 13327, MEMPHIS, TN 38113-0327
SINCE 1972

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling and the EPA registered labeling must be in the possession of the user at time of pesticide application.

All applicable directions, restrictions and use precautions on DREXEL SIMAZINE 4L label(s) are to be followed. This labeling contains supplemental instructions and restrictions for use of this product which may not appear on the DREXEL SIMAZINE 4L label. Follow the instructions carefully.

APPLICATION DIRECTIONS

Apply 1.6 to 2 pints of this product (0.8 to 1 lb. a.i.) per acre in the late Fall when the crop is well established. The application should be timed to coincide with the start of the Winter rainfall period to control germinating annual weeds. A second application at the rate of 1.6 to 2 pints of this product (0.8 to 1 lb. a.i.) per acre can be made in the Spring just after the last cultivation. Use a minimum of 20 gallons of spray solution with ground equipment per acre. For best results, field will have been cultivated one or more times and be relatively weed-free at time of treatment. Irrigation is not required to activate the herbicide if adequate rainfall is received.

USE RESTRICTIONS AND PRECAUTIONS:

- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.
- Do not use on crop, crop residue, or seed screenings for food or feed.
- Do not apply this product to sand, loamy sand, or gravelly areas of the field. Do not apply to areas of the field where water will accumulate.
- Do not rotate crops that are susceptible to simazine injury.
- For use under this SLN label, do not apply this product through any type of irrigation system.
- Aerial application is prohibited.

Special Crop Use Restrictions

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a nonfeed/nonfood use by the Oregon Department of Agriculture, and is regulated by the Oregon Administrative Rule (OAR) 603-057-0535, Pesticide Use On Crops Grown For Seed. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for this crop. Consequently, no portion of this seed may be used or distributed for food or feed. This restriction pertains to, but is not limited to, green chop, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage or seed screenings, and to the grazing of the crop field, stubble or regrowth. All seed screenings shall be disposed of in such a manner that screenings cannot be distributed or used for food or feed purposes, as indicated in OAR 603-057-0535. Additional regulations concerning seed screenings are stated in OAR 603-057-0535.

Any seed from the field treated with this product shall bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

“This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. This seed, in whole, as sprouts, or in any form may not be used for human consumption or animal feed. Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration, the Oregon Department of Agriculture and other regulatory agencies.”

This label is valid until December 31, 2012 or otherwise amended, withdrawn, cancelled, or suspended.