

Supplemental Label – Special Local Need Label

AgriSolutions DIURON 80DF (EPA Reg. No. 9779-318) EPA SLN OR-920023

For Preemergence And Postemergence Weed Control

THIS LABEL VALID UNTIL DECEMBER 31, 2009 OR UNTIL OTHERWISE AMENDED,
WITHDRAWN, CANCELLED, OR SUSPENDED.

FOR DISTRIBUTION AND USE ONLY WITHIN OREGON.

THIS LABEL AND THE MAIN LABEL MUST BE IN THE POSSESSION OF THE USER AT THE
TIME OF PESTICIDE APPLICATION.

ACTIVE INGREDIENT	
Diuron [3-(3,4-dichlorophenyl)-1,1-dimethylurea]	80%
INERT INGREDIENTS	20%
TOTAL	100%

CAUTION

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Sweep up spills and bury in a safe place away from water supplies.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves made of any waterproof material, and shoes plus socks.

FOR CONTROL OF CERTAIN BROADLEAF WEEDS AND ANNUAL GRASSES IN PERENNIAL RYEGRASS, TALL FESCUE, KENTUCKY BLUEGRASS, AND FINE FESCUE GROWN FOR SEED.

Apply this product only to well-established, vigorous stands of grasses as directed below. Use a sufficient volume of water, a minimum 25 gallons per acre, for thorough coverage of weed foliage. For best results, make application at the onset of fall rains and before weeds become established (typically October 1 through November 15). Weeds beyond the 2- to 4-leaf stage will usually not be controlled and it is desirable to remove them prior to applying this product. When making application, continue mechanical agitation of spray mixture and avoid overlapping spray patterns. Use higher rates within the range listed when treating larger weeds and heavier weed infestations. Weed control may be reduced where straw or ash residues have accumulated on the soil surface. Also, lack of moisture to activate the herbicide may reduce weed control. Tank mixtures or sequential treatment with other herbicides may reduce crop tolerance and increase risk of crop injury. When using this product in a tank mix or in a sequential treatment with other herbicides, do not use the maximum rates listed below unless compatibility and the potential for phytotoxicity have been evaluated. Crop tolerance may be reduced and the likelihood of crop injury may increase when crop is under stress caused by weather, diseases, insects and other such factors. Rotation to other crops within 2 years of application may cause injury to subsequent crops. Do not apply this product through any type of irrigation system.

Perennial Ryegrass (Established): Apply 1 to 2 pounds per acre per season (October 1 through mid-January) to control seedling grasses and broadleaf weeds such as annual bluegrass and others named on the product label.

Tall Fescue (Established): Apply 2 to 4 pounds per acre per season (October 1 through mid-January) to control seedling grasses and broadleaf weeds such as rattail fescue and others named on the product label.

Kentucky Bluegrass (Established stands east of the Cascade Mountains): Apply 1 _ to 3 pounds per acre per season (October 1 through mid-January) for suppression of rattail fescue and certain other seedling grasses and broadleaf weeds named on the product label. Downy brome is not controlled. Do not use on *Poa trivialis* grass seed varieties.

Fine Fescue (Illahee, Rainier, Pennlawn, Chewings and related varieties including hard fescue-Established stands west of the Cascade Mountains): Apply 1 to 2 pounds per acre for suppression of rattail fescue and certain other seedling grasses and broadleaf weeds named on the main product label. Make only 1 application per year. Do not use this product more than 2 years in succession on the same field.

Surface Water Protection Statement

Certain uses of diuron may be restricted by a U.S. District Court final order. You may refer to the Oregon Department of Agriculture web site at <http://oda.state.or.us/pesticide/lawsregs/buffers.html> or to the U.S. EPA web site at <http://www.epa.gov/espp/wtc/maps.htm#wtc6> for information regarding pesticides that may be impacted by the final order."