



**FITRA**

**Section 24(c) Special Local Need Label**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON**

**FUSILADE® DX Herbicide**

**EPA REG. NO. 100-1070  
EPA SLN NO. OR-950013**

**FOR CERTAIN FINE FESCUES GROWN FOR SEED USE ONLY**

This label valid until December 31, 2011 or until otherwise amended, withdrawn, cancelled, or suspended.

Active Ingredient:

Fluazifop-P-butyl

Butyl (R)-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoate\* ..... 24.5%

Other Ingredients ..... 75.5%

Total: ..... 100.0%

\*Fusilade DX herbicide contains 2 pounds (+) isomer (fluazifop-P-butyl) per gallon  
Contains petroleum hydrocarbons.

**Keep Out Of Reach Of Children**

**Caution**

## **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate, nitrile rubber, or neoprene rubber or viton
- Shoes plus socks

## **Environmental Hazards**

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area.

---

## **DIRECTIONS FOR USE**

---

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA-registered Fusilade DX Herbicide label and this label.

This label and the EPA-registered Fusilade DX Herbicide Section 3 label must be in the possession of the user at the time of application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## **Weeds Controlled**

For downy brome (cheatgrass) (*Bromus tectorum*), quackgrass (*Agropyron repens*) and bentgrass (*Agrostis sp.*) control in Creeping Red, Chewings and Hard Fescue varieties grown for seed in Oregon.

## **Application Instructions**

**Fall Application** - Apply FUSILADE DX at 8 oz. per acre to actively growing downy brome in the 2 to 4 inch stage. Apply at 12 to 16 oz. per acre to actively growing quackgrass or bentgrass in the 6 to 10 inch stage. Make a second application of 12 to 16 oz. per acre on the perennial grasses before they reach 10 inches of regrowth.

**Spring Application** - Apply FUSILADE DX at 8 to 16 oz. per acre to actively growing grasses. Downy brome in the 2 to 4 inch stage can be controlled by the lower rate but larger, established grasses (4 to 6 inches) may need the higher rate for adequate control. Use 12 to 16 oz. per acre in quackgrass and bentgrass in the 6 to 10 inch stage and a second application at the same rates before they reach 10 inches of regrowth.

## **RESTRICTIONS**

- **Do not apply FUSILADE DX after fescue grass seedhead develops into the boot stage.**
- **Treated fields may not be grazed or harvested for forage until after harvesting grass seed crop.**
- **Do not apply this product through any type of irrigation system.**

Fusilade® trademark of a Syngenta group company

24(c) registrant:  
Syngenta Crop Protection, Inc.  
P.O. Box 18300  
Greensboro, NC 27419-8300

EPA SLN No.: OR-950013

Label Code: OR1070063CA1006