



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

BEACON®

**EPA Reg. No. 100-705
EPA SLN No. OR-960025**

This label valid until December 31, 2012 or until otherwise amended, withdrawn, cancelled, or suspended.

**For Weed Control in Seedling and Established Stands of Kentucky
Bluegrass Grown for Seed Production Only**

Active Ingredient:

| | |
|---|------|
| Primisulfuron-methyl: 3-[4,6-Bis(difluoromethoxy)-pyrimidin-2-yl]-1-(2-methoxycarbonylphenyl-sulfonyl)urea: | 75% |
| Other Ingredients: | 25% |
| Total: | 100% |

Keep Out of Reach of Children

CAUTION

**SYNGENTA'S SPECIAL CONDITIONS, RISKS OF USE AND DISCLAIMER FOR
USE OF BEACON ON CROPS ON THIS 24C LABEL**

IMPORTANT- READ BEFORE USE

**THESE CONDITIONS RISKS OF USE AND DISCLAIMER ARE REQUIRED BY
SYNGENTA CROP PROTECTION INC AND NOT SPECIFIED BY U.S. EPA OR THE
STATE OF OREGON**

Labeling may be obtained through the Special Labels link at www.farmassist.com.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SYNGENTA CROP PROTECTION, INC. INTENDS THAT THE PRODUCT THAT IS THE SUBJECT OF THIS SECTION 24C LABEL BE PURCHASED ONLY BY END USERS WHO AGREE BY ELECTRONIC SIGNATURE ON SYNGENTA CROP PROTECTION'S INTERNET SITE TO THE TERMS AND CONDITIONS REQUIRED BY SYNGENTA CROP PROTECTION, INC. INCLUDING A WAIVER AND RELEASE FROM ALL LIABILITY AND INDEMNIFICATION BY THE USER AND/OR GROWER OF SYNGENTA AND OTHERS FOR FAILURE TO PERFORM AND FOR CROP INJURY, CROP YIELD REDUCTION, AND/OR CROP LOSS FROM USE OF BEACON ON CROPS ON THIS 24(C) LABEL. IF SUCH

TERMS AND CONDITIONS ARE UNACCEPTABLE, RETURN THE BEACON AT ONCE UNOPENED OR USE THE BEACON FOR A DIFFERENT APPROVED USE IN ACCORDANCE WITH THE LABEL AFFIXED TO THE PRODUCT CONTAINER.

USE OF BEACON (THE “PRODUCT”) ON CROPS LISTED (THE “CROP”) FOR THIS SPECIAL LOCAL NEED MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS AS FURTHER DISCUSSED BELOW. READ AND UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SPECIAL LOCAL NEED BEFORE USING THE PRODUCT ON THE CROP. SYNGENTA RECOMMENDS THAT THE USER TEST THIS PRODUCT TO DETERMINE ITS SUITABILITY FOR SUCH INTENDED USE.

Syngenta Crop Protection, Inc. makes the Product available for use in the manner described in this Supplemental Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury, Crop yield reduction or Crop loss. The decision to use this Product in the manner described in this Supplemental Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the potential risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any other relevant factors. Syngenta recommends that the user test this Product to determine its suitability for such intended use.

By purchasing the Product for use, or using the Product in the manner described in this Supplemental Labeling, you acknowledge and accept that, to the extent consistent with applicable law:

- 1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;
- 2) Syngenta Crop Protection, Inc. do not make, and do not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Supplemental labeling;
- 3) Syngenta Crop Protection, Inc. do not make, and do not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaim all warranties, expressed or implied, including any implied warranty of merchantability;
- 4) Syngenta Crop Protection, Inc. disclaim all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- 5) these conditions and Risks of Use for Special Local Need supersede any contrary representations or recommendations by Syngenta Crop Protection, Inc. or their respective agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Conditions and Risks of Use for Special Local Need are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

Environmental Hazards

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE

This label and the federal label for this product must be in the possession of the user at the time of pesticide application.

Follow all applicable directions, restrictions, and precautions on this Supplemental label and the main EPA-registered label. It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Refer to the EPA-registered Beacon label for additional weeds controlled and rotational crop restrictions.

This herbicide controls weeds by inhibiting a biochemical process which produces certain essential amino acids necessary for plant growth. The inhibited enzyme system is acetolactate synthase (ALS). Occurrence of ALS-resistant weed biotypes can be prevented or delayed by using this product in sequence with other herbicides having a different mode of action, and by using some form of mechanical control or a herbicide with a different mode of action to control weed escapes before they set seed.

Cleaning Equipment After Beacon Application

Because many crops are extremely sensitive to low rates of Beacon, special attention must be given to cleaning equipment before spraying a crop other than Kentucky bluegrass. Mix only as much spray solution as needed. Immediately after spraying, clean equipment thoroughly using this procedure.

1. Prepare a tank cleaning solution of 1 gal. of household ammonia per 50 gals. of water. Use sufficient cleaning solution to thoroughly rinse all surfaces.
2. When available, use a pressure rinser to clean the inside of the spray tank with this solution. Start agitation in the sprayer and thoroughly recirculate the cleaning solution for at least 15 minutes. All visible deposits must be removed from the spraying system.
3. Dispose of rinsate in an appropriate manner. Spray the cleaning solution on untreated Kentucky bluegrass, or return to a rinsate tank for later use as makeup water for spraying Kentucky bluegrass.
4. Repeat steps 1-3.
5. Remove nozzles and screens and clean separately after completing the above procedures to avoid injury to crops sprayed after the Beacon/Kentucky bluegrass application.

Seedling and Established Stands of Kentucky Bluegrass (*Poa pratensis* L.) Grown for Seed Production

Beacon can be applied postemergence to bluegrass with at least one tiller but prior to jointing for control of emerged weeds such as quackgrass, windgrass, partial control of cheat and downy brome, and the other weeds in Table 1. Beacon applications should be made to actively growing weeds. Kentucky bluegrass should be growing vigorously under favorable growing conditions.

Apply Beacon at 0.76 oz./A (one water-soluble packet for 2 acres). Alternatively, it may be applied as a split application of 0.38 oz./A (one-water soluble packet for 4 acres) followed by another 0.38 oz./A. If split application is used, the initial application should be made when weed sizes are within the range specified for optimum control in the following table; and the second half of the application should be made when the new weed growth or regrowth is again within the recommended size range for optimum control, generally 3-4 weeks after the initial application.

Application should be a broadcast spray with ground or aerial equipment. Refer to the EPA-registered Beacon label for mixing instructions, recommended spray volumes, and additive requirements. In general, a crop oil concentrate is the preferred additive when Beacon is applied alone for weed control in Kentucky bluegrass grown for seed. Tank-mixing Beacon with other herbicides is not recommended because of limited tank-mix testing to date; however, if Beacon is tank mixed with other products, the use of a nonionic surfactant instead of crop oil concentrate is recommended, to reduce the risk of crop injury from certain mixtures. **Syngenta will not be responsible for crop injury from tank mixtures of Beacon with other herbicides applied to spring or fall-planted seedlings of Kentucky bluegrass.** In addition, do not apply Beacon in tank mixture with, nor within 10 days before or 10 days after the application of, any organophosphate insecticide.

Aerial Application: Apply Beacon in water using a minimum spray volume of 3 gals. per acre. Include a nonionic surfactant, i.e., X-77®, at 1 qt. per 100 gals. of spray mix (0.25% volume/volume), or a good quality crop oil concentrate at no more than 2 pts. per acre (See Mixing Instructions on the EPA-registered Beacon label). Avoid application under conditions where uniform coverage cannot be obtained or where spray drift may occur. Make applications at a maximum height of 10 ft. above Kentucky bluegrass with low drift nozzles at a maximum pressure of 40 psi and wind speed not exceeding 10 mph to assure accurate Beacon application within the target area.

Restrictions/Precautions:

- **DO NOT** treat new seedlings of Kentucky bluegrass with Beacon if the plants have less than one well established tiller and are not in good vigor and growing condition. Applications can be made in the spring or fall, when the target weeds are in the recommended size range, and both the bluegrass and weeds are actively growing. **DO NOT** apply Beacon when the bluegrass is stressed due to drought, excessive heat, frost, flooding, poor fertility, diseases, insects, or other reason. Early applications to actively growing weeds and bluegrass give the best results.
- **DO NOT** irrigate within 4 hours after a Beacon application.
- **DO NOT** graze Beacon-treated Kentucky bluegrass fields within 90 days of application.
- **DO NOT** harvest within 60 days of application.
- **DO NOT** apply this product through any type of irrigation system.

- The total amount of Beacon applied during one crop season must not exceed 0.76 oz./A.
- Stunting of bluegrass plants may occur following application and, in some instances, seed yields may be adversely affected.

Beacon® trademark of a Syngenta Group Company

©2008 Syngenta

24(c) registrant:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

EPA SLN No.: OR-960025

Label Code: OR0705055CA0108