

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

This labeling and package must be in the possession of the user at the time of application. All applicable directions, restrictions and precautions on the package label and on this supplemental label must be followed.

WILDLIFE CONCERNS

To protect the Washington ground squirrel (*Spermophilus washingtoni*), use is prohibited in shrub-steppe and grassland habitat with deep loam and sandy-loam soils of northern Gilliam, Morrow, and Umatilla Counties. In addition, use is prohibited on any lands that lie within 250 meters of occupied Washington ground squirrel sites. For information regarding these restrictions and to determine if the desired location of application lies within the scope of this restriction, please call the Oregon Department of Fish and Wildlife at 541-676-5230.

Users are advised that the Migratory Bird Treaty Act makes it unlawful to kill any migratory bird except as permitted by regulation. **Any bird deaths caused by application of this pesticide is a violation of both state and federal law.** In North America, all birds are considered migratory except for upland game birds (e.g. grouse, ptarmigans, prairie-chickens, quail, ring-necked pheasant and wild turkey) and introduced bird species (e.g. European starling, house sparrow, and rock dove). Refer to full product label for Endangered Species Considerations.

USE RESTRICTIONS: For control of voles (*Microtus spp.*), house mice, deer mice, and certain ground squirrels in grasses grown for seed as specified below.

- **APPLICATIONS ARE ONLY ALLOWED FROM MAY 8 THROUGH AUGUST 31. DO NOT APPLY FROM SEPTEMBER 1 THROUGH MAY 7.**
- Do not make applications if geese or other migratory birds have been observed in the field within the last seven (7) days.
- If geese are observed in the treated field within seven (7) days of the application, hazing is recommended. A hazing permit is not required
- Do not apply this product by air.
- This product cannot be injected or introduced into any type of irrigation system (e.g. chemigation).

RESTRICTED USE PESTICIDE DUE TO HAZARDS TO NONTARGET SPECIES

For retail sale and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

ZP[®] RODENT BAIT AG

EPA Reg. No. 12455-17

EPA SLN No. OR-990034

ACTIVE INGREDIENT:

Zinc Phosphide (CAS #1314-84-7) 2%

INERT INGREDIENTS:..... 98%

TOTAL 100%

SPECIAL LOCAL NEED SUPPLEMENTAL LABEL

FOR CONTROL OF VOLES, MICE AND GROUND SQUIRRELS IN GRASSES GROWN FOR SEED AND ONLY FOR USES AND APPLICATION METHODS ON THIS SUPPLEMENTAL LABEL

For Distribution and Use Only in the State of Oregon.

This label is valid until April 30, 2010 or until otherwise amended, withdrawn, canceled, or suspended.

KEEP OUT OF REACH OF CHILDREN CAUTION

(See full product label for additional precautionary statements)

24 (c) Registrant:



Bell Laboratories, Inc.
3699 Kinsman Blvd.
Madison, WI 53704

USE RESTRICTIONS (Continued)

- Do not graze animals on treated area or harvest any part of the treated crop for livestock feed for forty five (45) days after application.
- Do not exceed (4) applications per treated area per year.
- Do not apply if rain is forecast within the next 24 hours.
- Anyone who applies bait by ground equipment must wear a dust mask and chemically resistant gloves. Applicators must wear chemically resistant gloves when hand baiting.

APPLICATION DIRECTIONS (Directions for three methods provided):

PESTS: For control of the vole complex (*Microtus spp.*); house mouse (*Mus musculus*); and deer mouse (*Peromyscus maniculatus*).

Preharvest - Broadcast

When all restrictions indicated on this label are met, this product may be broadcast by ground driven dispensing devices, cyclone seeder or by hand. Apply at a rate of 6 - 10 pounds per acre. Use the higher label rate in areas with highest active population level. The number of vole holes is not an indicator of population level, or activity.

Do not broadcast bait on bare ground.

Post Harvest – Between Windrows

Between windrow use is allowed at or after swathing. Apply to areas of high infestation between windrows. Apply with a ground driven dispensing device (i.e. attached to combine), cyclone spreader or by hand. Apply at a rate of 6-10 pounds per acre. Only apply between windrows. Do not apply over, or on windrows. Do not apply in such a manner that results in zinc phosphide bait to be combined. Incinerate or compost seed screenings.

After treating areas between windrows:

- Do not bale straw for any purpose including, stack burning, disposal, animal feed or bedding.
- Do not remove straw from the field.
- Do not use any portion of the crop, including seed screenings or straw for animal feed or bedding.

Application Directions (Continued next page)

APPLICATION DIRECTIONS (Continued)

Post Harvest – Broadcast Use

Post-harvest broadcast use is only allowed after all seed/plant by-products (including straw) have been removed from field, or after seed is removed and straw-load is left in field. This product may be applied with a ground driven dispensing device, cyclone spreader or by hand. Apply at a rate of 6-10 pounds per acre. After treatment of the field, do not remove straw from the field; or bale straw for any purpose including, stack burning, disposal, animal feed or bedding. Applications may not be made after August 31.

IMPROVING EFFICACY

- Avoid applying zinc phosphide at rates less than allowed by this label. Voles/mice may develop a learned aversion to zinc phosphide when exposed to sub-lethal rates.
- Zinc phosphide bait deteriorates when exposed to rain, humidity and wet acidic soils. The longer the product is exposed to these conditions, the less effective it will be. Efficacy may be improved if applications are made after the soil pH is increased. Avoid treating before morning dew evaporates.

APPLICATION DIRECTIONS:

Spot Baiting Use (allowed May 8 – August 31)

PESTS: For control of California ground squirrel (*Spermophilus beechyi*); Columbian ground squirrel (*S. columbianus*); and Townsend ground squirrel (*S. townsendii*).

Prebaiting: To increase acceptance of treated bait by ground squirrel; prebait with one teaspoon of untreated grain within a 6 inch diameter around each active burrow 2-3 days prior to using toxic bait.

Handbaiting: Apply one teaspoon of bait within a 6 inch diameter around each active burrow. Do not place in piles.

24 (c) Registrant:



Bell Laboratories, Inc.
3699 Kinsman Blvd.
Madison, WI 53704