



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

Bravo Weather Stik[®]
EPA REG. NO. 50534-188-100
EPA SLN NO. OR-990040

For Control of Downy Mildew on Sugarbeets Grown for Seed

Active Ingredient:

Chlorothalonil (tetrachloroisophthalonitrile).....	54.0%
Other Ingredients:	46.0%
Total	100.0%

Contains 6.0 Pounds Chlorothalonil Per Gallon (720 grams per liter)

This label valid until December 31, 2012 or until otherwise amended, withdrawn, canceled, or suspended.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Environmental Hazards

This product is toxic to aquatic invertebrates and wildlife. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This label and the federal label for this product must be in the possession of the user at the time of pesticide application.

Follow all applicable directions, restrictions, and precautions on this Supplemental label and the main EPA-registered label. It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, or pets either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

CROP	DISEASE	Pts. Product./A (lbs. a.i./A)	APPLICATION DIRECTIONS
Sugarbeets (seed production only)	Downy mildew	1.7 (1.3)	Use in sufficient water to obtain adequate coverage, normally 20 to 150 gallons per acre for dilute sprays and 5 to 10 gallons per acre for concentrate ground sprays and aircraft applications. Make the first application in mid-October when disease conditions become favorable, and make additional applications as favorable disease conditions warrant, through April of the following year. Do not apply Bravo Weather Stik to sugar beet plants after bolting and seedhead emergence has occurred. Bravo Weather Stik may be applied through sprinkler irrigation systems. See <i>Application and Calibration Techniques for Chemigation</i> on Bravo Weather Stik label.

Special Crop Use Restrictions

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a nonfeed/non-food use by the Oregon Department of Agriculture, and is regulated by Oregon Administrative Rule (OAR) 603-057-0535, Pesticide Use On Crops Grown For Seed. If the applicator of this pesticide is not the producer, the applicator should provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, should provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for this crop. Consequently, no portion of this seed crop may be used or distributed for food or feed for 1 year (365 days) after the last application of this product. This restriction pertains to, but is not limited to: green chop, forage, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage or seed screenings, and to the grazing of the crop field, stubble or regrowth. All seed screenings shall be disposed of in such a manner that the screenings cannot be distributed or used for food or feed purposes, as indicated in OAR 603-057-0535. Additional regulations concerning seed screenings are stated in OAR 603-057-0535.

Any seed from a field treated with this pesticide product shall bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

"This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. This seed, in whole, as sprouts, or in any form, may not be used for human consumption or animal feed. Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration, the Oregon Department of Agriculture and other regulatory agencies."

Certain uses of chlorothalonil may be restricted by a U.S. District Court final order. You may refer to the Oregon Department of Agriculture web site at <http://egov.oregon.gov/ODA/PEST/buffers.shtml> or to the U.S. EPA web site at <http://www.epa.gov/espp/wtc/maps.htm#wtc6> or <http://www.epa.gov/espp/wtc/uselimitation.htm> for information regarding pesticides that may be impacted by the final order.

Bravo Weather Stik[®] is a trademark of a Syngenta Group Company

Section 24(c) Registrant:
G.B. Biosciences, Inc.
Greensboro, NC 27409

Section 24(c) Distributor:
Syngenta Crop Protection, Inc.
Greensboro, NC 27409

OR0188061EA0807