



Section 24(c) Special Local Need Label

GB BIOSCIENCES CORPORATION

Syngenta Crop Protection, Inc.
(Agent for GB Biosciences)
PO Box 18300
Greensboro, NC 27419-8300
Telephone: 336-632-6000

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

DACONIL ULTREX®

EPA Reg. No. 50534-202
EPA SLN No. WA-000003

FOR CONTROL OF ORNAMENTAL BULB DISEASES

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL
MAY RESULT IN POOR PEST CONTROL OR CROP INJURY.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA-registered Daconil Ultrex label.

This label must be in the possession of the user at the time of application.

DISEASES	RATE PER 100 GALLONS	APPLICATION DIRECTIONS
Basal rot Neck rot Other bulb rots including those caused by Mucor, Zygorrhynchus, Rhizopus, Curvularia and Rhizoctonia	2.5-5.0 lbs	Dip bulbs from 15 minutes up to 4 hours prior to planting. Add recommended DACONIL ULTREX dilute mixture to dip tanks to maintain dip solution at levels needed to achieve complete bulb coverage. Wear chemical-resistant gloves during application and when handling treated bulbs.

Do not apply this product to ornamental bulbs through any type of irrigation system. Refer to the EPA registered label for DACONIL ULTREX, EPA Reg. No. 50534-202, for other directions for use.

This product is toxic to aquatic invertebrates and wildlife. DO NOT apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. DO NOT contaminate water when disposing of equipment wash water or rinsate. Daconil Ultrex should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service or US Fish & Wildlife Service for information on listed threatened or endangered species (e.g. Bull trout, Chinook salmon). Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

This chemical can contaminate surface water through spray drift. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow ground water, areas with infield canals or ditches that drain to surface waters, areas not separate from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

APPLICATION OF DIP-TANK TREATMENT WATER: Spent dip-tank treatment water may be applied using ground equipment to bulb fields for basal, neck or other bulb rots under the following conditions:

- The concentration of Daconil Ultrex in solutions to be applied to bulb fields must not exceed 5% Daconil Ultrex (5 lb. of Daconil Ultrex per 100 gallons water).
- Do not apply more than a total of 36.4 lbs. active ingredient of chlorothalonil (i.e. 44 lbs. Daconil Ultrex) per acre per year as a dip-tank solution to bulb fields or as a spray to field grown bulb plants during the growing season.
- Apply only when rainfall is not expected for at least 24 hours after application.
- Do not apply Daconil Ultrex using ground equipment within 25 feet of surface water.

This label for DACONIL ULTREX expires and must not be distributed or used in accordance with this SLN registration after December 31, 2007.

DACONIL ULTREX® is a trademark of Syngenta Crop Protection, Inc.

24(c) registrant: GB Bioscience Corp.
Greensboro, NC 27419-8300

EPA SLN No.: WA-000003

WADaconilUltrexOrnamentalBulbsSLNLbl-GBB 1-24-03