



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

DACONIL ULTREX®

EPA REG. NO. 50534-202-100
EPA SLN NO. WA-000003b

For Control of Ornamental Bulb Diseases

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL OR CROP INJURY.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

This labeling must be in the possession of the user at the time of application.

DISEASES	RATE PER 100 GALLONS	APPLICATION DIRECTIONS
Basal rot Neck rot Other bulb rots including those caused by Mucor, Zygorrhynchus, Rhizopus, Curvularia and Rhizoctonia	2.5-5.0 lbs	Dip bulbs from 15 minutes up to 4 hours prior to planting. Add recommended DACONIL ULTREX dilute mixture to dip tanks to maintain dip solution at levels needed to achieve complete bulb coverage. Wear chemical-resistant gloves during application and when handling treated bulbs.

Restrictions/Precautions

Do not apply this product to ornamental bulbs through any type of irrigation system. Refer to the EPA registered label for DACONIL ULTREX, EPA Reg. No. 50534-202-100, for other directions for use.

This product is toxic to aquatic invertebrates and wildlife. DO NOT apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. DO NOT contaminate water when disposing of equipment wash water or rinsate. Daconil Ultrex should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program web site at

<http://agr.wa.gov/PestFert/EnvResources/EndangSpecies.htm>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered.

This chemical can contaminate surface water through spray drift. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow ground water, areas with infield canals or ditches that drain to surface waters, areas not separate from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

Certain uses of chlorothalonil may be restricted by a U.S. District Court final order. You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/EnvResources/Buffers.htm> for information regarding pesticides that are impacted by the final order.

APPLICATION OF DIP-TANK TREATMENT WATER: Spent dip-tank treatment water may be applied using ground equipment to bulb fields for basal, neck or other bulb rots under the following conditions:

- The concentration of Daconil Ultrex in solutions to be applied to bulb fields must not exceed 5% Daconil Ultrex (5 lb. of Daconil Ultrex per 100 gallons water).
- Do not apply more than a total of 36.4 lbs. active ingredient of chlorothalonil (i.e. 44 lbs. Daconil Ultrex) per acre per year as a dip-tank solution to bulb fields or as a spray to field grown bulb plants during the growing season.
- Apply only when rainfall is not expected for at least 24 hours after application.
- Do not apply Daconil Ultrex using ground equipment within 25 feet of surface water.

WSDA Container Disposal Guidance: Completely empty the contents of the pesticide container into application equipment, and apply to a labeled crop or site. The empty container may be disposed of in a sanitary landfill. Burning is not a legal method of container disposal in Washington.

This label for DACONIL ULTREX expires and must not be distributed or used in accordance with this SLN registration after December 31, 2011.

DACONIL ULTREX® is a trademark of Syngenta Crop Protection, Inc.

24(c) Sub-registrant:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

EPA SLN No.: WA-000003b

WA0202042CA0707