



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

Betamix® Herbicide
EPA Reg. No. 264-621
For Control of Ladysthumb, Pale smartweed,
Wild mustard, Shepherd's-purse,
Hedgemustard, Pigweeds (*Amaranthus* spp.),
Hairy Nightshade, and Common Chickweed in
Table Beets and Swiss Chard Grown for Seed.
For Use Only by Members of the Puget Sound
Seed Growers Association
EPA SLN. No. WA-010015

24(c) Supplemental Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

This label for Betamix® Herbicide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2009.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Failure to follow the Directions for Use and Precautions on this label may result in poor product performance, crop injury, or illegal residues.

Apply Betamix® Herbicide as a postemergence broadcast spray to table beets or Swiss chard grown for seed using ground application equipment in 20 to 50 gallons of water (see Chart 1 for broadcast rates). A second Betamix® Herbicide treatment may be applied 4 weeks after the first application. (See complete Betamix® Herbicide label for recommendations concerning conditions that may cause stress, spray equipment, and cultural practices).

**CHART 1
Dosage Chart for Broadcast Application of Betamix® Herbicide**

Weed Stage	Betamix® Herbicide (Pints/Acre)
2 leaf*	1.5 – 2.3
> 3 leaf	1.5 – 3.0

* Do not use the 3 pint rate when plants are at the 2 leaf stage.

Tank Mixes: The weed control spectrum may be improved by tanking mixing Betamix® Herbicide with Nortron® SC Herbicide in 20 to 50 gallons of water (see Chart 2 for broadcast rates).

**CHART 2
Dosage Chart for Tank Mixes of Betamix® Herbicide and Nortron® SC Herbicide**

Weed Stage	Pints/Acre Broadcast	
	Betamix® Herbicide	Nortron® SC Herbicide
4 – 5 leaf	2.0	1.0
6 – 8 leaf	3.0	2.0

* Do not use the 3 pint rate when plants are at the 2 leaf stage.

RESTRICTIONS / PRECAUTIONS:

- Do not apply Betamix® Herbicide after the crop begins to form buds.
- For use under this SLN label, do not apply this product through any type of irrigation system.
- For use under this SLN label, do not apply this product using aerial equipment.
- This pesticide is toxic to fish and aquatic organisms. Betamix Herbicide should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WADA Endangered Species Program web site at <http://agr.wa.gov/PestFert/EnvResources/EndangSpecies.htm>, or contact the Washington Department of Fish & Wildlife, National Marine fisheries Service (NOAA Fisheries) or U.S. Fish & Wildlife Service for information regarding aquatic species listed a threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.
- All table beet and Swiss chard seed screenings shall be disposed of in such a way that they cannot be distributed or used for food or feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the WSDA forthwith upon request. Disposal records shall consist of documentation from a controlled dumpsite, incinerator, or other equivalent disposal site and shall show the lot numbers, amount of material disposed of, its grower(s), and the date of disposal. No portion of the table beet and Swiss chard seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes. Table beet and Swiss chard seed shall bear a tag or container label, which forbids use of the seed for human consumption or animal feed. Table beet and Swiss chard seed may not be distributed for human consumption or animal feed. Producers of table beet or Swiss chard grown for seed who use this product are required to inform, in writing, conditioners receiving seed produced on fields treated with this product. A copy of this labeling is required to be provided to the conditioners by the producer. At the processing plant, processed seed must be labeled "Not for human or animal consumption".

CONTAINER DISPOSAL GUIDANCE: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the internet from WSU at <http://pep.wsu.edu/waste/wd.html> or from WSDA at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

WAIVER OF LIABILITY

Bayer CropScience intends that this Section 24(c) label be distributed only by the Puget Sound Seed Growers Association only to end users and/or growers who agree in writing to the terms and conditions required by the Puget Sound Seed Growers Association including a waiver and release from all liability and indemnification by the user and/or grower of Bayer CropScience, Puget Sound Seed Growers Association, and others for failure to perform and crop damage from the use of Betamix® Herbicide on table beets or Swiss chard grown for seed. If such terms and conditions are unacceptable, return Betamix® Herbicide at once unopened.

This product when used on table beets or Swiss chard grown for seed may lead to crop injury, loss or damage. Bayer CropScience recommends that the user and/or grower test this product in order to determine its suitability for such intended use. The Puget Sound Seed Growers Association and Bayer CropScience make this product available to the user and/or grower solely to the extent the benefit and utility, in the sole opinion of the user and/or grower, outweigh the extent of potential injury associated with the use of this product. The decision to use or not to use this Betamix® Herbicide must be made by each individual Betamix® Herbicide user and/or grower on the basis of possible crop injury from Betamix® Herbicide, the severity of weed infestation, the cost of alternative (pest) controls, and other factors. Bayer CropScience intends that because of the risk of failure to perform or crop damage that all such use is at the user's and/or grower's risk.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at *www.bayercropscienceus.com*