



MANZATE® 75DF FUNGICIDE  
SUPPLEMENTAL LABELING

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

MANZATE® 75DF FUNGICIDE

EPA Reg. No. 1812-414  
EPA SLN No. WA-010018

FOR DISEASE CONTROL ON  
SMALL SEEDED VEGETABLE SEED CROPS

Directions for Use:

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Application Instructions:

Begin applications when disease first appears or is reported in the area. Apply 2 pounds of Manzate® 75DF Fungicide per acre in thorough coverage sprays. Repeat applications on a 7 to 10 day application schedule until the threat of disease is over.

CROP	DISEASES CONTROLLED
<i>Allium</i> spp. (Leek, bunching and dry bulb onions)	Botrytis Leaf Blight ( <i>Botrytis</i> spp.) Downy Mildew ( <i>Peronospora destructor</i> ) Neck Rot ( <i>Botrytis</i> spp.) Purple Blotch ( <i>Alternaria porri</i> )
Arugula	Alternaria Leaf Spot ( <i>Alternaria</i> sp.)
Beet	Alternaria Leaf Spot ( <i>Alternaria</i> sp.) Cercospora Leaf Spot ( <i>Cercospora beticola</i> ) Downy Mildew ( <i>Peronospora farinosa</i> ) Rust ( <i>Uromyces betae</i> )
Carrot	Leaf Blight ( <i>Alternaria dauci</i> ) Leaf Spot ( <i>Cercospora carotae</i> ) Rust ( <i>Uromyces lineolatus</i> )
Coriander	Leaf Spot ( <i>Alternaria</i> spp.)
Crucifers (Brassica and Raphanus spp.)	Black Spot ( <i>Alternaria alaternata</i> and <i>A. brassicola</i> ) Cercospora Leaf Spot ( <i>Cercospora brassicola</i> ) Downy Mildew ( <i>Peronospora parasitica</i> )
Broccoli Raab, Brussels Sprouts, Cabbage, Cauliflower, Chinese Cabbage, Chinese Mustard, Collard, Kale, Kohlrabi, Rutabaga, Turnip, Mustard, Radish, Rape and Spinach	
Dill	Alternaria Leaf Spot ( <i>Alternaria</i> sp.)
Endive	Downey Mildew ( <i>Bremia lactucae</i> )
Lettuce	Anthrachnose ( <i>Microdochium panattonianum</i> ) Cercospora Leaf Spot ( <i>Cercospora longissima</i> ) Downy Mildew ( <i>Bremia lactucae</i> )
Parsley	Alternaria Leaf Spot ( <i>Alternaria</i> sp.)
Parsnip	Alternaria Leaf Spot ( <i>Alternaria</i> sp.)

CROP	DISEASES CONTROLLED
Spinach	Downy Mildew ( <i>Peronospora effusa</i> and <i>P. farinosa</i> ) Leaf Spot ( <i>Alternaria</i> spp.) Leaf Spot ( <i>Cercospora beticola</i> ) Leaf Spot ( <i>Cladosporium variable</i> )
Swiss Chard	Cercosora Leaf Spot ( <i>Cercospora beticola</i> ) Rust ( <i>Uromyces betae</i> )

**RESTRICTIONS/PRECAUTIONS:**

All screenings from seed crops listed on this SLN label shall be disposed of in such a way that they cannot be distributed or used for food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the director immediately upon request. Conditioner disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, composter or other equivalent disposal site and shall include lot numbers, amounts of material disposed of, the grower(s), and the date of disposal.

No portion of the plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings, may be used or distributed for food or feed purposes.

Seed crops listed on this SLN shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.

Seed crops listed on this SLN may not be distributed for human consumption or animal feed.

For use under this SLN supplemental label, DO NOT apply this product through any type of irrigation system.

Apply no more than 12 applications per 12 month period.

**NOTE:**

This labeling contains new or supplemental instructions for use of this product which may not appear on the package label. Follow instructions carefully.

**SPRAY DRIFT MANAGEMENT:** A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, and relative humidity) and method of application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

**WSDA Container Disposal Guidance:**

Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the internet from WSU at <http://pep.wsu.edu/waste/wd.html> or from WSDA at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

**This label for Manzate® 75DF expires and must not be distributed or used in accordance with this SLN registration after December 31, 2011.**

**GSL-678 020807 021307**

Griffin L.L.C.  
Stine-Haskell Research Center  
P.O. Box 30  
Newark, DE 19714-0030