



SIPCAM AGRO USA, INC.

**For Distribution and Use Only within the state of Washington
For Control of Ascochyta Blight in Chickpeas**

Echo[®] 720 Agricultural Fungicide

EPA Reg. No. 60063-7

EPA SLN No. WA-020012

- It is a violation of Federal law to use this product in a manner that is inconsistent with its labeling.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- This labeling must be in the possession of the user at the time of pesticide application.

DIRECTIONS FOR USE

Application Rates: Dosage rates on this label indicate pints of ECHO 720 per acre, unless otherwise stated. Under conditions favoring disease development, the high rate specified and shortest application interval should be used. For chickpeas, the listed maximum individual and seasonal application rates must not be exceeded and the listed minimum retreatment intervals must not be decreased.

DISEASES	PHI (DAYS)	RATE PER APPLICATION PER ACRE	APPLICATION DIRECTIONS FOR CHICKPEAS
Ascochyta blight	14	1-3 / 8 to 2 pints	For use only on chickpeas (dry beans) to be harvested dry with pods removed. Use sufficient water to obtain thorough coverage. Begin application at first onset of disease, which can occur 2 to 4 weeks before flowering. Repeat applications at 7 to 10 day intervals. DO NOT apply more than five (5) times per growing season. Echo 720 Agricultural Fungicide may be applied through sprinkler equipment. Refer to the EPA registered label for chemigation directions. RESTRICTIONS: Do NOT allow livestock to graze in treated areas or feed treated plant parts to livestock. Do NOT apply more than a total of 6.0 lbs chlorothalonil active ingredient (i.e. 1 gallon of Echo 720 Agricultural Fungicide) per acre per year.

This product is toxic to aquatic invertebrates and wildlife. Echo 720 Agricultural Fungicide should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/EnvResources/EndangSpecies.htm>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered species. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

Certain uses of chlorothalonil may be restricted by a U.S. District Court final order. You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/EnvResources/Buffers.htm> for information regarding pesticides that are impacted by the final order.

WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

This label for Echo 720 Agricultural Fungicide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2012.

ECHO is a registered trademark of Sipcam Agro USA, Inc.

24(c) Registrant:
Sipcam Agro USA, Inc.
300 Colonial Center Parkway, Suite 230
Roswell, Georgia 30076

11/2/07