



PIPERA

Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

SWITCH® 62.5WG

**EPA Reg. No. 100-953
EPA SLN No. WA-020016**

For control of Alternaria Leaf/pod spot (*Alternaria brassicicola* and *A. brassicae*), white mold (*Sclerotinia sclerotiorum*) and gray mold (*Botrytis cinerea*) on selected brassica vegetable crops grown for seed

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA-registered Switch 62.5WG label.

This labeling must be in the possession of the user at the time of herbicide application.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

BRASSICA GROWN FOR SEED

Broccoli, Chinese Broccoli, Broccoli raab, Brussels sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Kale, Chinese Kale, Kohlrabi, Mustard Greens, Chinese Mustard, Mustard Spinach, Radish (other than Daikon), Rape Greens, Rutabaga; Turnip, Turnip Greens.

Application Rate

Apply 14 oz. of Switch 62.5WG (0.219 pounds fludioxonil + 0.328 pounds cyprodinil) per acre.

Application Method/Frequency/Timing

Apply in sufficient water to provide thorough coverage using ground application equipment only. Begin application preventively at early bloom and repeat on 7 to 10 day intervals depending on disease pressure. Begin applications prior to or at the onset of disease and repeat application on a 7-10 day interval if conditions remain favorable for disease development.

Resistance Management

After 2 applications of Switch 62.5WG, alternate with another fungicide with a different mode of action for 2 applications.

Restriction/Precautions

- Switch 62.5WG may be applied up to 7 days prior to harvest (i.e., 7-day PHI).
- Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.
- Do not apply more than 56 oz/A of Switch 62.5WG per crop/year
- Do not apply more than 1.3 lb a.i./A of cyprodinil containing products and 0.9 lb a.i./A of fludioxonil containing products per crop/year

Seed Crop Restrictions

1. All brassica seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the director immediately upon request. Conditioner disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, or composter, other equivalent disposal site and shall include the lot numbers, amount of material disposed of, the grower(s), and the date of disposal.
2. No portion of the brassica seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
3. Brassica seed shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.
4. Brassica seed may not be distributed for human consumption or animal feed.

Producers of brassica seed who use this product are required to inform, in writing, conditioners receiving seed produced on fields treated with this product. A copy of this labeling is required to be provided to the conditioner by the producer.

Surface Water Advisory

This chemical may contaminate water through drift of spray in wind. This chemical has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this chemical. A level, well maintained vegetative buffer strip between areas to which this chemical is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this chemical will be reduced by avoiding applications when conditions favor runoff (such as when soils are saturated and/or significant rainfall is forecast in the next 48 hours). Sound erosion control practices will reduce this chemical's contribution to surface water contamination.

Aquatic Toxicity

This pesticide is toxic to fish, aquatic invertebrates, oysters, and shrimp. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

Switch 62.5WG should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the Washington State Department of Agriculture Endangered Species Program website at <http://agr.wa.gov/PestFert/NatResources/EndangSpecies.htm>, or contact the Washington Department of Fish and Wildlife, national Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult the federal label or additional restrictions and precautions to protect aquatic organisms.

WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington."

Chemigation: For use under this SLN label do not apply this product through any type of irrigation system.

This label for Switch 62.5WG expires and must not be distributed or used in accordance with this SLN registration after December 31, 2013.

Switch® trademark of a Syngenta Group Company

© 2008 Syngenta

24(c) registrant:
Syngenta Crop Protection, Inc.
P.O. Box 18300, Greensboro, NC 27419-8300

WA0953038DA0408