

RESTRICTED USE PESTICIDE

Due to acute toxicity

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

Guthion® Solupak 50% Wettable Powder Crop Insecticide

For Use as a Border Treatment Around Alkali Bee Beds to Control Black Oil Beetle.

EPA Reg. No. 264-733

SLN No. WA-030025

24(c) Supplemental Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

Guthion® Solupak 50% Wettable Powder Crop Insecticide must not be sold or distributed for use under this SLN after December 31, 2010. This label expires and must not be used after September 30, 2012.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

SITE	INSECT	METHODS AND RATE OF APPLICATION
Border treatment around alkali bee beds	Black oil beetle (<i>Meloe niger</i>)	Ground Spray – Initial applications should be made prior to emergence of overwintering beetles. Apply Guthion Solupak 50% Wettable Powder Crop Insecticide at the rate of 1 lb in 8 gallons of water per 4,000 feet. Apply no more than 20 lbs per acre per application in sufficient water to ensure complete coverage of the area around but not on the bee bed. Repeat applications as necessary at 4 to 5 day intervals during the time beetles are emerging to maintain control. Pitfall traps for spot treatment - Install edging along bee beds and sprinkler 1-3 tablespoons (0.5 - 1.5 ounces) of Guthion Solupak 50% Wettable Powder Crop Insecticide per large pitfall trap or buried container (e.g. five-gallon plastic bucket) prior to emergence of beetles (e.g. early March). Add sufficient water into the trap at the end of season and apply spent mixture into the ground depression left after the removal of the trap.

RESTRICTIONS/PRECAUTIONS

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or broadleaf weeds. Do not apply this product or allow it to drift to blooming crops or broadleaf weeds.

For use under this SLN label, do not apply this product through any type of irrigation system.

This pesticide is extremely toxic to fish and wildlife. Guthion Solupak 50% Wettable Powder Insecticide should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/EnvResources/EndangSpecies.htm>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

WSDA Container Disposal Guidance: Completely empty the contents of the pesticide container into application equipment, and apply to a labeled crop or site. The empty container may be disposed of in a sanitary landfill. Burning is not a legal method of container disposal in Washington.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).