

RESTRICTED USE PESTICIDE

Due to Reproductive Effects

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification. Direct supervision for this product is defined as the certified applicator being physically present during mixing, loading, equipment repair and equipment cleaning. Certified applicators must ensure that all persons involved in these activities under their direct supervision are informed of the precautionary statements.

REGISTRATION FOR SPECIAL LOCAL NEED FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

MSR™ SPRAY CONCENTRATE

SYSTEMIC INSECTICIDE FOR EFFECTIVE SYSTEMIC CROP PROTECTION

EPA Reg. No. 10163-220

EPA SLN NO.: WA-060005

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA-registered label.
- This labeling must be in the possession of the user at the time of application.

CROP	PEST	PINTS	COMMENTS
Newly planted and nursery-grown Non-Bearing Caneberries	Aphids (<i>including green peach aphid</i>)	1 – 1 ½	Apply specified dosage before canopy closure as a full coverage spray in sufficient water for complete coverage.
<ul style="list-style-type: none">• Do not apply more than once per acre per season.• Do not apply within 365 days of harvest.• Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 16 days.• Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.• For use under this SLN label do not apply this product through any type of irrigation system or by air.• A no-spray buffer zone of 25 feet (groundboom) is required between the application site and any area managed for wildlife or wildlife habitat.			

WSDA CONTAINER DISPOSAL GUIDANCE: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the internet from WSU at <http://pep.wsu.edu/waste/wd.html> or from WSDA at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and wildlife.

MSR Spray Concentrate should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service or US Fish & Wildlife Service for information on listed threatened or endangered species (e.g. Bull trout, Chinook salmon). Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

IMPORTANT: This product is sold subject to the Conditions of Sale set forth on the container label.

This label for MSR Spray Concentrate expires and must not be distributed or used in accordance with this SLN registration after December 31, 2009

24(c) REGISTRANT

Gowan Company
P.O. Box 5569
Yuma, AZ 85366

WA-060005 MSR nonbearing cane (revised 4-17-06)