



Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

OSPREY™ Herbicide
EPA Reg No. 264-802

**For Control of Italian Ryegrass and Wild Oats
 in Fall-Sown or Winter Triticale.**

EPA SLN No. WA-060010

24(c) Supplemental Label

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF
 WASHINGTON**

This label for Osprey™ Herbicide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2009.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

WEED APPLICATION TIMING

OSPREY™ Herbicide is a postemergent herbicide with best results being obtained when applications are made to young actively growing weeds. For control or suppression of labeled grasses, the weed application timing is from 1-leaf to the 2-tiller stage of growth. See weed tables for appropriate application timing and weed size. Treat heavy weed infestations before they become competitive with the crop.

SPRAY ADDITIVES

OSPREY™ Herbicide is a water dispersible granule that does not include an adjuvant. A recommended adjuvant **must** be tank mixed with OSPREY™ Herbicide according to the guidelines as described in the Mixing Order section of the Federal label.

APPLICATION METHODS

Ground Application

OSPREY™ Herbicide can be applied broadcast in 10 or more gallons of water per acre. For weed control in dense weed canopies, use 15 or more gallons of water per acre. Weed infestations should be treated before they become competitive with the crop.

Aerial Application

Calibrate the spray equipment prior to use. OSPREY™ Herbicide should be applied in a minimum of 5 gallons of water per broadcast acre. Damage to sensitive crops can occur as a result of spray drift. Spray drift can be managed by several application factors and by spraying under the appropriate climatic conditions. The use of nozzles and spray pressure that deliver MEDIUM spray droplets as indicated in the nozzle manufacturer's catalogs and in accordance with ASAE Standard S-572 are highly recommended for optimum spray coverage and canopy penetration. DO NOT use raindrop nozzles. Aerial applications with this product should be made at a maximum height of 10 feet above the crop with low drift nozzles at a maximum pressure of 40 psi. Avoid application under conditions where uniform coverage cannot be obtained.

WEEDS CONTROLLED WITH OSPREY™ HERBICIDE ON FALL-SOWN OR WINTER TRITICALE

Product	Rate	Weed	Crop
OSPREY™ Herbicide	3.2 ounces/acre	Wild oat (<i>Avena fatua</i>) Windgrass (<i>Aperi spica-venti</i> & <i>Apera interrupta</i>)	Apply OSPREY™ Herbicide from emergence up to the jointing stage of triticale development.
	4.75 ounces/acre	Annual (Italian) ryegrass (<i>Lolium multiflorum</i>) Annual bluegrass (<i>Poa annua</i>)	

WEEDS SUPPRESSED WITH OSPREY™ HERBICIDE ON FALL-SOWN OR WINTER TRITICALE

Product	Rate	Weed	Crop
OSPREY™ Herbicide	4.75 ounces/acre	Downy brome (<i>Bromus tectorum</i>) Jointed goatgrass (<i>Aegilops cylindrical</i>) Ripgut brome (<i>Bromus rigidus</i>) Hairy chess (<i>Bromus commutatus</i>)	Apply OSPREY™ Herbicide from emergence up to the jointing stage of triticale development.

RESTRICTIONS/PRECAUTIONS FOR USE

- For use under this SLN label do not apply this product through any type of irrigation system.
- Do not apply under circumstances where possible drift to unprotected persons or to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption can occur.
- All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.
- For ground applications, the applicator must:
 1. Apply when there is sustained wind away from native plant communities, OR
 2. Use low-pressure nozzles according to manufacturer's specifications that produce only coarse or very coarse droplets, OR
 3. Leave 50 foot untreated buffer between treatment area and native plant communities.
- For aerial applications:
 1. The distance of the outer most nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
 2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.
 3. The applicator must apply when there is sustained wind away from native plant communities or leave a 350-foot untreated buffer between treatment area and native plant communities.
- A recommended adjuvant must be tank-mixed with OSPREY™ Herbicide according to the guidelines on the federal label.
- Applications should be made to young, actively growing weeds. Control may be reduced when weeds are under stress due to severe weather conditions, drought, very cold temperatures, etc.
- Do not apply within 30 days of harvesting triticale forage, and 60 days for hay, grain and straw.
- Follow all applicable use directions, weed and crop application timings, precautions and restrictions on the product label for OSPREY™ Herbicide.

Container Disposal Guidance: Completely empty the contents of the paper container into application equipment, apply to a labeled crop or site. The empty container may be disposed in a sanitary landfill. Burning is not a legal method of container disposal in Washington.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).