

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON



Supplemental
Labeling

(Prowl® H₂O herbicide contains 3.8 pounds of pendimethalin per gallon formulated as an aqueous capsule suspension)

EPA Reg. No. 241-418

EPA SLN No.: WA-060018

FOR WEED CONTROL IN FIRST YEAR NON-BEARING STRAWBERRIES

NOTICE

BASF CORPORATION ("BASF") HAS OBTAINED A REGISTRATION AS A SERVICE TO THOSE GROWERS WHO WISH TO USE PROWL H₂O HERBICIDE AS AN AID IN CONTROLLING WEEDS IN FIRST YEAR NON-BEARING STRAWBERRIES. BASF DOES NOT WARRANT THE USE OF PROWL ON FIRST YEAR NON-BEARING STRAWBERRIES. THE DECISION TO APPLY THIS HERBICIDE MUST BE MADE BY THE GROWER WITH THE UNDERSTANDING THAT REDUCED CROP GROWTH, QUALITY, YIELD AND/OR DELAYED MATURITY MAY RESULT EVEN WHEN DIRECTIONS FOR USE ARE FOLLOWED.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of application.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered product label for Prowl® H₂O herbicide, EPA Reg. No. 241-418.

DIRECTIONS FOR USE

Observe all cautions and limitations on this label and on the labels of products used in combination with **Prowl H₂O**. Do not use **Prowl H₂O** other than in accordance with the instructions set forth on this label. The use of **Prowl H₂O** not consistent with this label may result in injury to crops. Keep containers closed to avoid spills and contamination.

Read the **Precautionary Statement, Environmental Hazards, Storage and Disposal, and Disclaimer** statements appearing on the container label.

GENERAL INFORMATION

Prowl H₂O may be applied for pre-emergence weed control of most annual grasses and certain broadleaf weeds as they germinate, but it will not control established weeds. Unusually cold, excessively wet, or hot and dry conditions that

delay germination or extend it over a long period of time may reduce weed control.

Over-application can result in crop stand loss, crop injury, or rotational crop injury.

Uneven application or improper soil incorporation can decrease weed control or cause crop injury. Soil incorporation deeper than recommended can reduce weed control.

Use of **Prowl® H₂O herbicide** in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of this product and, therefore, rotational crop injury is always possible.

Refer to the **Prowl H₂O** container label for a complete listing of weeds controlled.

APPLICATION INSTRUCTIONS

Stunting, reduced growth, or reduction in daughter plants may occur with this use. Apply **Prowl H₂O** preemergence prior to transplanting strawberries. Make one application in 10 or more gallons of water per acre with ground equipment.

APPLICATION RATES

Broadcast Rate Per Acre of Prowl H₂O Used In First Year Non-Bearing Strawberries (Pints Per Acre)

Soil Texture	<3% Organic Matter	>3% Organic Matter
Coarse	1.5 – 2.0 pts.	2.0 – 3.0 pts.
Medium	2.0 – 2.5 pts.	2.0 – 3.0 pts.
Fine	2.0 – 3.0 pts.	2.5 – 3.5 pts.

RESTRICTIONS AND LIMITATIONS

For use under this SLN label, **DO NOT** apply this product through any type of irrigation system or by air.

DO NOT use on peat or muck soils.

Prowl H₂O is most effective when incorporated in the weed germination zone by rainfall or irrigation.

DO NOT exceed the maximum labeled rate for a given soil texture.

DO NOT plant winter wheat or winter barley in the fall following **Prowl H₂O** application at rates above 3.0 pints per acre.

NOTE: If loss of strawberry stand occurs due to adverse weather conditions, any crop registered for **Prowl H₂O** preplant incorporated use can be replanted the same year into **Prowl H₂O** treated soil without adverse effects. If replanting is necessary, soil should not be reworked deeper than the **Prowl H₂O** treated zone

DO NOT harvest for food or feed any portion of the strawberry plant within 1 year (365 days) of **Prowl H₂O** application.

ENDANGERED SPECIES

This pesticide is toxic to fish. **Prowl H₂O herbicide** should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/EnvResources/EndangSpecies.htm>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage: **Prowl H₂O herbicide** freezes around 15°F and is stable under conditions of freezing and thawing. Product that has been frozen should be thawed and recirculated prior to use.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law.

WSDA Container Disposal Guidance:

Plastic Containers: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the Internet from WSU at <http://pep.wsu.edu/waste/wd.html> or from WSDA at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county.

This label for Prowl[®] H₂O herbicide expires and must not be distributed or used in accordance with this FIFRA 24(c) Special Local Need registration after December 31, 2011.

©2006 BASF Corporation. All rights reserved
Prowl is a registered trademark of BASF.

0000241-00418.20051018e.NVA 2005-04-195-0289

24(c) Registrant:

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709


The Chemical Company