



Bayer CropScience LP  
 P.O. Box 12014  
 2 T.W. Alexander Drive  
 Research Triangle Park, North Carolina 27709  
 1-866-99BAYER (1-866-992-2937)  
<http://www.bayercropscienceus.com>

42-S Thiram Fungicide  
 For Use On Parsley Seed (For Export to Mexico Only)  
 EPA Reg. No. 264-929  
 EPA SLN. No. WA-060023

## 24(c) Supplemental Label

### FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

CROP	DISEASE	DOSAGE
Parsley Seed (For Export to Mexico Only)	To aid in the control of seed decay, damping-off and seedling blight	For standard mechanical, slurry or mist-type treaters, apply 2.5 to 5 fluid ounces of 42-S Thiram Fungicide per 100 pounds of seed. Product should be diluted with sufficient water to give uniform coverage of seed.
NOTE: Treat and conduct germination tests on a small portion of the seed before committing the total seed lot to the chemical treatment.		
SEED LABELING: Treated seed must be labeled "For Export Only – Not to be sold or offered for sale in the United States". The U.S. Seed Act requires that treated seed be labeled with the following information: <p style="margin-left: 40px;">This seed treated with Thiram at the rate recommended by the manufacturer. Treated seed must not be used for, or mixed with food or animal feed or processed for oil. Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water.</p> <p>Use with an EPA-approved dye that imparts an unnatural color to the seed.</p>		

FOR EXPORT ONLY TO MEXICO. This label is based solely on treatment requirements made by Mexico requiring Thiram be applied to imported parsley seed. Bayer CropScience makes no claims as to the effect of this product on disease control or seed germination. All such risks shall be assumed by the Buyer.

WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the internet from WSU at '<http://pep.wsu.edu/waste/wd.html>' or from WSDA at '<http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>'. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

This label for 42-S Thiram Fungicide must not be distributed or used in accordance with this SLN registration after December 31, 2011.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at [www.bayercropscienceus.com](http://www.bayercropscienceus.com)