

Focus™ SC

EPA Reg. No. 10163-275

§ 24(c) Special Local Need Registration SLN No. WA-070003 FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

For Control of Powdery Mildew on Cherries

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Focus SC should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program website at <http://agr.wa.gov/Pest/Fert/EnvResources/EndangSpecies.htm>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries for information regarding aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Focus SC¹

Crop Disease Spray Schedule	Use rate per acre (fl oz)	Use rate per 100 gal. of water dilute ² (fl oz)	Maximum spray interval (days)	Use Directions and Restrictions
Cherries Powdery Mildew (late bloom-harvest)	8-12	2-3	14-21	<p>Application timing: Focus SC may be applied alone or in tank mix combination with recommended rates of a protectant fungicide. Begin applications at delayed dormant and continue as needed until harvest.</p> <p>Restrictions: Do not apply more than 12 fl. oz. of Focus SC per acre per application. Use the higher rate range for severe infection levels. Do not apply more than two (2) consecutive sprays of Focus SC or another DMI (e.g. Rubigan) fungicide. Do not apply more than 36 fl. oz. or three (3) applications of Focus SC per acre per year. Focus SC may be applied up to "harvest". For use under this SLN do not apply this product through any type of irrigation system. Apply Focus SC with ground equipment only.</p>

¹Dilute sprays are sprayed until point of runoff. Gallonage of dilute sprays will vary depending on tree size, density of canopy and stage of growth of the trees and tree spacing in the orchard.

WSDA CONTAINER DISPOSAL GUIDANCE

Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop of site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the internet from WSU at <http://agr.wa.gov/PestFert/EnvResources/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

This label for Focus SC expires and must not be distributed or used in accordance with this SLN registration after December 31, 2011.

24(c) registrant:
Gowan Company
PO Box 5569
Yuma, AZ 85366-5569

EPA SLN No.:
Effective Date:
Revised:

WA-070003
2/16/07