

syngenta

FIFRA

Section 24(c) Special Local Need Label

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF
WASHINGTON**

RIDOMIL GOLD[®] SL

**EPA Reg. No. 100-1202
EPA SLN No. WA-070012**

**For Control Of Cavity Spot Caused By
Pythium spp. In Carrots**

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL
MAY RESULT IN POOR PEST CONTROL, ILLEGAL RESIDUES, OR CROP INJURY.

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This label must be in the possession of the user at the time of pesticide application.

Follow all applicable directions, restrictions, and precautions including statements pertaining to
the Worker Protection Standards, on the EPA-registered Ridomil Gold SL label.

DIRECTIONS FOR USE - Carrots

Control of Cavity Spot caused by *Pythium* spp.

Soil Applications

Pre-plant incorporated: Apply Ridomil Gold SL at 0.5 to 1.3 pts./A per treated acre as a broadcast soil application in sufficient water or liquid fertilizer to provide uniform coverage and incorporate in the top two inches of soil. Use a minimum carrier volume of 10 gallons/acre.

At planting surface applications: Apply Ridomil Gold SL at 0.5 to 1.3 pts/A immediately after planting or prior to emergence in sufficient water or liquid fertilizer to provide uniform coverage and incorporate in with irrigation. Use a minimum carrier volume of 10 gallons/acre. For chemigation applications prior to carrot emergence, apply 4 to 8 ounces per acre in sufficient water to provide uniform coverage and incorporate in the irrigation water. In sandy soils, apply a maximum of ½ inch of irrigation after the chemigation event. Refer to the Ridomil Gold SL label (EPA Reg. No. 100-1202) for use directions for chemigation. This product may be applied through irrigation systems. Refer to the EPA registered label for chemigation instructions.

Banded applications: Apply Ridomil Gold SL in a 7-inch band. Use the formula in the General Information section of the Ridomil Gold SL label to calculate the amount of Ridomil Gold SL needed per acre. Use a minimum carrier volume of 10 gallons/acre. If natural rainfall is not expected before the seeds start germinating, Ridomil Gold SL should be incorporated mechanically before planting or be moved into the seed zone after planting with irrigation.

Post-Plant Applications:

Beginning 28-50 days after planting, apply Ridomil Gold SL on a 14 to 21 day interval. Apply 0.25 to 1 pint of Ridomil Gold SL per acre in water by chemigation, by ground equipment with a spray directed to the base of the plant or shanked in with liquid fertilizer. Use a minimum carrier volume of 10 gallons/acre. All ground applications must be followed by irrigation with at least one inch of water to properly move the product into the root zone. **DO NOT** exceed 1.5 pints per acre on post plant applications.

Notes: For best control of cavity spot caused by *Pythium* spp. use a preventive disease control program that incorporates an at planting or seed treatment use of Ridomil Gold SL or Apron XL followed by one or more supplemental applications

Restrictions

- To avoid illegal residues, do not apply more than 2.8 pts. of Ridomil Gold SL per acre per crop season.
- If a seed treatment containing mefenoxam or metalaxyl is used, do not use an at-planting soil application.
- Do not exceed the equivalent of 0.65 lb ai/A per crop of soil-applied and 0.75 lb ai/A per crop of post-planting applied mefenoxam containing products.

Pre-harvest Interval

DO NOT harvest carrots within 7 days of a Ridomil Gold SL application or illegal residues may result.

WSDA Container Disposal Guidance

Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the internet from WSU at <http://pep.wsu.edu/waste/wd.html> or from WSDA at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

This label for Ridomil Gold SL expires and must not be distributed or used in accordance with this SLN registration after December 31, 2009.

Ridomil Gold® trademark of a Syngenta Group Company

24(c) registrant:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

Label Code: WA1202092AA0707