


RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

EPA SLN NO.
WA-070017



FOR DISTRIBUTION AND
USE ONLY WITHIN THE
STATE OF WASHINGTON

Brigade[®] 2EC

Insecticide/Miticide

EPA REG. No. 279-3313

EMERGENCY CALLS: 800-331-3148

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARD REQUIREMENTS AND PRECAUTIONS ON THE EPA REGISTERED LABEL.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.

Crop	Pest	Rate of Application
Carrot Grown for Seed (Hybrid or Open Pollinated) Parsley Grown for Seed Parsnip Grown for Seed	Aphids Armyworms Cutworms Loopers Lygus Bugs Spider Mites	3.9 to 6.4 ounces per acre

Directions for Use on:

Apply Brigade 2EC at the rate of 3.9 to 6.4 ounces (0.06 to 0.1 lb. active) per acre. Apply in a minimum of 8 gallons of water per acre by air equipment or in a minimum of 20 gallons of water by ground equipment.

Restrictions / Precautions:

For use under this SLN label do not apply this product through any type of irrigation system.

Do not apply more than 6.4 ounces (0.1 lb. active) per acre per application. Do not make more than 2 applications per season. Maintain a minimum of 21 days between applications. Rotational crops may be planted no sooner than 30 days after last application.

Do not apply by ground equipment within 25 feet, or by air within 150 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.

Do not cultivate within 10 feet of aquatic area so as to allow growth of a vegetative filter strip.

This pesticide is extremely toxic to fish and aquatic invertebrates. Brigade 2EC should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may



FMC Corporation
Agriculture Products Group
1735 Market Street
Philadelphia, PA 19103

Brigade and FMC -FMC Trademarks
©2007 FMC Corporation

refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/EnvResources/EndangSpecies.htm>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

All Carrot, Parsley and Parsnip seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the director immediately upon request. Conditioner disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, composter or other equivalent disposal site and shall include the lot numbers, amounts of material disposed of, the grower(s), and the date of disposal.

No portion of the Carrot, Parsley and Parsnip seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.

Carrot, Parsley and Parsnip seed shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.

Carrot, Parsley and Parsnip seed may not be distributed for human consumption or animal feed.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply the 3.9 to 6.4 oz/acre (0.06 to 0.1 lb ai/acre) rate of this product to blooming Carrot, Parsley and Parsnip grown for seed: apply as a pre-bloom or post-bloom spray only. Applications of the 2 oz/acre (0.032 lb ai/acre) rate of this product to blooming Carrot, Parsley and Parsnip grown for seed must be timed to coincide with periods of minimum bee activity, between late evening and midnight.

Note: Information is provided on use of the 2 ounce rate during bloom in order to minimize the hazard to pollinators. While use of the 2 ounce rate is legal, FMC does not recommend the 2 ounce rate, since efficacy may not be as good as the full label rate. Users wishing to apply the low rate during bloom must be willing to accept reduced product efficacy, in order to protect pollinators.

WSDA Container Disposal Guidance:

Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

THIS LABEL FOR BRIGADE 2EC EXPIRES AND MUST NOT BE DISTRIBUTED OR USED IN ACCORDANCE WITH THIS SLN REGISTRATION AFTER DECEMBER 31, 2012.



FMC Corporation
Agriculture Products Group
1735 Market Street
Philadelphia, PA 19103

Brigade and  -FMC Trademarks
©2007 FMC Corporation