



Arysta LifeScience

FIFRA 24(c) REGISTRATION
EPA SLN No. WA-080002
Expiration Date: 12/31/2013

REGISTRATION FOR SPECIAL LOCAL NEED
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

IPRODIONE 4L AG

EPA Reg. No. 66330-297

FOR USE ON BRASSICA CROPS GROWN FOR SEED

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

CROP: Brassica crops grown for seed (Broccoli raab, Brussels sprouts, Cabbage, Cauliflower, Chinese cabbage, Chinese kale, Chinese mustard, Collard, Kale, Kohlrabi, Mustard, Radish (other than daikon), Rape, Rutabaga, Spinach mustard, Turnip).

TARGET PEST: Alternaria leaf and pod blight, Sclerotinia white rot.

RATE: Use 2 to 4 pints of product per acre per application. The higher rate should be used when severe disease conditions exist.

FREQUENCY AND TIMING OF APPLICATION: Application should be made at full bloom, at pod set, and just prior to harvest if favorable disease conditions exist.

DILUTION RATE AND METHOD OF APPLICATION: Apply in 20 to 100 gallons of water per acre by ground. Apply in a minimum of 10 gallons of water per acre by air. Addition of a suitable resin-based sticker is recommended according to dilution rates on the sticker label. Thorough coverage is necessary for effective control.

This product may be applied through irrigation systems. Refer to the EPA registered label for chemigation directions.

WSDA Chemigation Guidance:

- Application off-site is prohibited. The chemigation application must be continuously observed whenever sensitive areas as defined in WAC 16-202-1002(44) (including but not limited to schools, parks, dwellings, occupied buildings or structures, public roadways, and waters of the state) are at risk of being exposed to drift, runoff, or overspray. In order to minimize the potential for application off-site, WSDA recommends that the product only be applied through low pressure irrigation systems (defined as 2 to 35 pounds/square inch) with a nozzle release height no higher than 3 feet above the target crop, and that end guns be disabled throughout the application.
- An inspection port or a direct access point is required, and it must be positioned immediately upstream of the irrigation mainline check valve and be of sufficient size to allow visual and manual inspection of the check valve and low pressure drain. The inspection port or access point must have a minimum diameter of four inches, if feasible (WAC 16-202-1012[1]).
- The chemigation application tank cannot be placed within 20 feet of the wellhead or other sensitive areas. Mixing or loading activities cannot occur within 20 feet of the wellhead or other sensitive areas (WAC 16-202-1008[1]).
- WSDA Chemigation Rules (WAC 16-202-1001 through WAC 16-202-1024), and information on USEPA Authorized Alternative Chemigation Safety Equipment,

Distribution Uniformity and other chemigation topics are available on the WSDA website (<http://agr.wa.gov/PestFert/ChemFert/default.htm>).

RESTRICTIONS / PRECAUTIONS:

- Iprodione 4L AG should not be applied to blooming Brassica crops grown for seed during the pollination period, due to concerns with toxicity to developing honey bee larvae.
- All Brassica crop seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the WSDA immediately upon request. Disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, composter or other equivalent disposal site and shall include the lot numbers, amount of material disposed of, the grower(s), and the date of disposal.
- No portion of the Brassica seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
- Brassica seed crop shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.
- Brassica seed crop may not be distributed for human consumption or animal feed.
- Iprodione 4L AG is a dicarboximide fungicide. Repeated use of dicarboximide fungicides may result in resistance to Iprodione.
- This pesticide is toxic to invertebrates. Iprodione 4L AG should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/NatResources/EndangSpecies.htm>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.
- If applying this product adjacent to a water body such as a lake, reservoir, river, permanent stream, marsh or natural pond, estuary, or commercial fish pond, there must be at least a 25-foot vegetative buffer strip between the water body and the point of application.
- This pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitats for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

WSDA Container Disposal Guidance:

Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

This label for Iprodione 4L AG expires and must not be distributed or used in accordance with this SLN registration after December 31, 2013.

Manufactured For:
Arysta LifeScience North America Corporation
15401 Weston Parkway #150
Cary, NC 27513