



FIFRA

Section 24(c) Special Local Need Label

RESTRICTED USE PESTICIDE
Due to Toxicity in Fish, Mammals, and Aquatic Organisms

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

ZORO® Miticide/Insecticide

EPA Reg. No. 67760-71
EPA SLN No. WA-080003

For control of twospotted spider mites on alfalfa grown for seed

DIRECTIONS FOR USE

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
 - Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.
 - This label must be in the possession of the user at the time of application.
-

Alfalfa Grown For Seed

Apply ZORO® when twospotted spider mites first appear. **DO NOT** apply more than 32 fluid ounces or more than two (2) applications of ZORO® per acre in any growing season. If a second application is necessary for control, do not apply within 21 days of the first application and only after an alternative miticide with a different mode of action has been used.

Ground application: Apply 8-16 fluid ounces of product (0.01-0.02 pounds of active ingredient) per acre by ground application in a minimum of 20 gallons of water per acre.

Aerial application: Apply 12-16 fluid ounces of product (0.014-0.02 pounds of active ingredient) per acre by aerial application in a minimum of 10 gallons of water per acre.

Restrictions/Precautions:

- Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.
- For use under this SLN label, do not apply this product through any type of irrigation system.

- This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming alfalfa grown for seed if bees are visiting the treatment area. Applications must be timed to coincide with periods of minimum bee activity, between late evening and midnight.
- All alfalfa seed screenings shall be disposed of in such a way that they cannot be distributed or used for food or feed. The seed conditioner shall keep records of screening disposal for three years from the date of the disposal and shall furnish the records to the WSDA immediately upon request. Disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, composter or other equivalent disposal site and shall include the lot numbers, amount of material disposed of, the grower(s), and the date of disposal.
- No portion of the alfalfa seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
- Alfalfa seed shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.
- Alfalfa seed may not be distributed for human consumption or animal feed.
- This pesticide is toxic to fish and wildlife. ZORO® should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/NatResources/EndangSpecies.htm> or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.
- Do not apply by ground within 25 feet, or by air within 150 feet of lakes, reservoirs, rivers, permanent streams, marshes, pot holes, or natural ponds, estuaries and commercial fish farm ponds.
- Do not cultivate within 25 feet of the aquatic area so as to allow growth of a vegetative filter strip.

WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

Expiration Date: This label for ZORO® expires and must not be distributed or used in accordance with this SLN registration after December 31, 2013.

24(c) registrant:
 Cheminova, Inc.
 1700 Route 23, Suite 300
 Wayne, NJ 07470
www.cheminova.us.com

ZORO® is a registered trademark of Cheminova

Rev. 04/22/2008