



Arysta LifeScience

FIFRA 24(c) REGISTRATION  
EPA SLN No. WA-080005  
Expiration Date: 12/31/2013

REGISTRATION FOR SPECIAL LOCAL NEED  
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

## **IPRODIONE 4L AG**

EPA Reg. No. 66330-297

### **FOR COMMERCIAL SEED TREATMENT OF BRASSICA CROPS GROWN FOR SEED**

#### **DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

**CROP/SITE:** Brassica crops grown for seed: Arugula, Broccoli, Broccoli raab, Brussels sprouts, Cabbage, Cauliflower, Chinese broccoli, Chinese cabbage, Collard, Kale, Kohlrabi, Mustard, Radish (other than daikon), Rape, Rutabaga, Spinach mustard, and Turnip.

**TARGET PEST:** Black leg (*Phoma lingam*).

#### **RATE/METHOD OF APPLICATION:**

**Seed Treatment:** Apply as a seed soak. Use 8 fluid ounces of Iprodione 4L AG in 6 gallons of water to treat 3 lbs. of Brassica seed for 24 hours at 30° C. Allow the treated seeds to dry thoroughly before packaging or planting. Do not apply more than 1 application to the seeds.

**Slurry Treatment:** Use 16 fluid ounces of Iprodione 4L AG per 100 lbs of Brassica seed as a slurry with enough water to insure complete coating of the seed. Allow the treated seeds to dry thoroughly before packaging or planting. Do not apply more than 1 application to the seeds.

#### **RESTRICTIONS/PRECAUTIONS:**

- For use in commercial seed treatment facilities only. Do not apply on farm.
- Brassica seed treated with this product shall only be used for planting Brassica crops grown for seed production.
- Bags containing treated seed shall be labeled with the following: "This seed has been treated with Iprodione 4L AG (iprodione). Do not use for feed, food, or oil purposes. Not for human or animal consumption. Store away from feeds and foodstuffs."
- All Brassica seed treated with this product containing iprodione must be colored with an EPA-approved dye (e.g. 40 CFR 180) which imparts an unnatural color to the seed to prevent the inadvertent use of treated seed for food, feed, or oil purposes.
- Crop injury may result under unfavorable environmental conditions.
- All Brassica crop seed screenings shall be disposed of in such a way that they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the WSDA immediately upon request. Disposal records shall consist of documentation of on-farm disposal, disposal at a controlled dumpsite, incinerator, composter or other equivalent disposal site and shall include the lot numbers, amount of material disposed of, the grower(s), and the date of disposal.

- No portion of the Brassica seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
- Brassica seed crop shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.
- Brassica seed crop may not be distributed for human consumption or animal feed.
- This pesticide is toxic to invertebrates. Iprodione 4L Ag should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/NatResources/EndangSpecies.htm>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

**WSDA Container Disposal Guidance:**

Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

This label for Iprodione 4L Ag expires and must not be distributed or used in accordance with this SLN registration after December 31, 2013.

Manufactured For:  
Arysta LifeScience North America Corporation  
15401 Weston Parkway #150  
Cary, NC 27513

Rev. 04-23-2008