



**FIFRA 24(c) SUPPLEMENTAL LABELING**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON**

**ETHEPHON 2**

EPA Reg. No. 66330-262

EPA SLN No. WA-080009

**FOR USE ON DOUGLAS FIR SEED ORCHARDS**

- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

**DIRECTIONS FOR USE**

A foliar spray of Ethephon 2 can be used to chemically eradicate unwanted seed cones that may become habitat for crop damaging insects.

**DOUGLAS FIR SEED ORCHARDS**

<b>Crop</b>	<b>Rate</b>	<b>Spray Volume Gallons/ Acre</b>	<b>Specific Directions</b>	<b>Remarks</b>
Douglas fir seed orchards	2 2/3 pints per 100 gallons of water	100 - 500	Apply as a foliar spray to Douglas fir seed cones when upright and the scales are open.  Use sufficient water for thorough coverage. Spray volume is dependent on the sprayer and tree size.	Best results are obtained when Ethephon 2 is used in combination with a nonionic surfactant, such as Induce <sup>®</sup> , at a rate of 4 pints/100 gallons.  Efficacy declines as seed cone scales close and begin to turn pendent.

**RESTRICTIONS / PRECAUTIONS**

- Do not enter or allow worker entry during the restricted entry interval (REI) of 48 hours. The REI is 72 hours in areas where average rainfall is less than 25 inches per year.
- For use under this SLN registration do not apply this product through any type of irrigation system.
- Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.
- This pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).



Arysta LifeScience

**WSDA Container Disposal Guidance:**

- Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA web site at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

**This label for Ethephon 2 expires and must not be distributed or used in accordance with this SLN registration after December 31, 2013.**

24(c) registrant:

Arysta LifeScience North America Corporation  
15401 Weston Parkway Suite150  
Cary, NC 27513

Induce<sup>®</sup> is a registered trademark of Helena Chemical Company

Rev: 8/4/2008