



FIFRA 24(c) REGISTRATION

**REGISTRATION FOR SPECIAL LOCAL NEED
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON**

**EVITO 480 SC Fungicide
EPA REG. NO. 66330-64 EPA SLN No: WA-080011**

FOR USE ON POTATO (BANDED AND IN-FURROW APPLICATIONS)

DIRECTIONS FOR USE

- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

POTATO

Soilborne / Seedling Diseases Controlled	Rate to Use	Application Timing
Black scurf (<i>Rhizoctonia solani</i>)	0.16 – 0.24 fl oz./1000 row feet	See directions and rates under SOILBORNE / SEEDLING DISEASE CONTROL.

SOILBORNE / SEEDLING DISEASE CONTROL

Evito can provide control of soilborne diseases, including Black scurf, if applied early in the growing season. Specific applications for soilborne diseases include in-furrow applications or banded applications applied over the row, either shortly after plant emergence or during herbicide applications or cultivation. These applications will provide control of pre- or post-emergence damping off and diseases that infect plants at the soil-plant interface. The use of either type of application depends on the cultural practices in the region. In some locations, one type of application may provide better disease control than the other, depending on the timing of the disease epidemic. Seedling diseases are generally controlled by in-furrow applications while banded applications are more effective against soilborne diseases that develop later in the season. Consult your local expert for guidance regarding application type.

Banded applications: Apply Evito prior to infection as a directed spray to the soil, using single or multiple nozzles, adjusted to provide thorough coverage of the lower stems and the soil surface surrounding the plants. Band width should be limited to 7 inches or less. Apply Evito at a rate of 0.16-0.24 fl. oz. product/1000 row feet. These applications come into contact with the foliage and are counted as foliar applications when considering resistance management. They may be applied during cultivation or hilling operations to provide soil incorporation.

In-furrow applications: Apply Evito as an in-furrow spray in 3-7 gallons of water at planting. Mount the spray nozzle so the spray is directed into the furrow just before the seed are covered. Use the higher rate when the weather conditions are expected to be conducive for disease development, if the field has a history of *Pythium* problems, or if minimum/low till programs are in place.

IN-FURROW APPLICATION RATES

RATE PER 1000 ROW FEET	PRODUCT PER ACRE (fl. oz.)						
	22" rows	30" rows	32" rows	34" rows	36" rows	38" rows	40" rows
fl. oz. product							
0.16	3.8	2.8	2.6	2.5	2.3	2.2	2.1
0.24	-	-	3.9	3.7	3.5	3.3	3.1

40" = 13,068 row ft., 38" = 13,754 row ft., 36" = 14,520 row ft., 34" = 15,374 row ft., 32" = 16,315 row ft., 30" = 17,424 row ft., and 22" = 23,760 row ft./Acre

RESTRICTIONS / PRECAUTIONS:

- Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.
- Do not apply more than 22.8 fl oz (0.72 lbs a.i.) of EVITO 480 SC per acre per year (including banded and in-furrow applications).
- There is a maximum number of 6 applications per season, and a minimum interval of 7 days between applications.
- Do not use the 0.24 fl. oz/1000 row feet rate on 22 or 30 inch rows.
- Do not apply EVITO 480 SC within 7 days of harvest.
- For use under this SLN label do not apply this product through any type of irrigation system.
- This pesticide is toxic to fish and aquatic invertebrates. Evito 480 SC should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/NatResources/EndangSpecies.htm>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

WSDA CONTAINER DISPOSAL GUIDANCE:

Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

EVITO is a trademark of Arysta LifeScience North America Corporation

This label for EVITO 480 SC expires and must not be distributed or used in accordance with this SLN registration after December 31, 2013.

24(c) Registrant:

Arysta LifeScience North America Corporation
15401 Weston Parkway, Suite 150
Cary, NC 27513

Rev. 08-29-2008