

## RESTRICTED USE PESTICIDE

Due to high acute toxicity and toxicity to wildlife.

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



**Bayer CropScience**

Bayer CropScience LP  
P.O. Box 12014  
2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)

Nemacur® 3 Emulsifiable Systemic Insecticide-Nematicide

For Use on Iris and Narcissus Bulbs to Control Nematodes.

EPA Reg. No. 264-731

SLN. No. WA-090006

### *24(c) Supplemental Label*

## FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

Sale and distribution of existing stocks of Nemacur 3 Emulsifiable Systemic Insecticide-Nematicide by persons other than Bayer CropScience may continue until March 31, 2009. This SLN label expires and must not be used after December 31, 2010.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

CROP	PEST	DOSAGE NEMACUR® 3	REMARKS
Iris and Narcissus Bulbs	Nematodes	2 to 3-1/3 gallons per acre (12,445 linear feet of row)	Apply specified dosage in the furrow at time of planting, so that the material is incorporated into the covering soil, or in a 10- to 12-inch band over the row after planting. Apply in furrow in the fall at time of planting or in a band over the row in October, November or December. <b>NOTE:</b> On year old plantings, apply over the row as a fall application in October, November or December. Incorporate with irrigation or rainfall as soon as possible after application. Tolerance of bulbs to NEMACUR may vary with varieties. Test a small number of plants if tolerance is not locally known. Do not apply to green foliage.

### RESTRICTIONS

- Do not apply this product with knapsack or similar equipment that is placed on the user's body. Do not use mist sprayers or aerial spray equipment.
- Do not make more than one application per crop season.
- Do not apply more than 3-1/3 gallons (10 lb ai fenamiphos) per acre per year.
- Do not apply within 100 feet of the following aquatic areas: lakes, reservoirs, rivers, permanent streams, marshes, natural ponds and estuaries.
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.
- Do not apply to hydrologic soil group A soils that are excessively drained and predominately sand or loamy sand with water tables less than 50 feet deep. These classifications and soil taxonomy refer to USDA definitions.
- For use under this SLN label do not apply this product through any type of irrigation system.

**GROUND WATER ADVISORY:** Fenamiphos is known to leach through soil and has been found in ground water as a result of agriculture use. Users are advised not to apply in areas where soils are permeable, particularly where ground water is used for drinking water.

This pesticide is toxic to fish and wildlife. NemaCur® 3 Emulsifiable Systemic Insecticide-Nematicide should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may contact the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/EnvResources/EndangSpecies.aspx>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

**WSDA Container Disposal Guidance:** Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.aspx>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at [www.bayercropscienceus.com](http://www.bayercropscienceus.com)

NemaCur is a registered trademark of Bayer.