

Product Bulletin



Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON
ORTHENE® 75 S Soluble Powder (Water Soluble Packets)

FOR AN APPLICATION OF ORTHENE 75 S SOLUBLE POWDER (WATER SOLUBLE PACKETS)
WITH A MINIMUM OF 75 DAYS TO HARVEST (POST-BLOOM) CRANBERRIES
FOR CONTROL OF BLACKHEADED FIREWORM

EPA Reg. No. 59639-26
EPA SLN No. WA-960024
Rev. 3

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of application.

Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA-registered label.

Crop	Insects*	Amount <i>Orthene 75 S</i> (WSP) Per	Use Directions/Restrictions	Days to Harvest
Cranberries	Blackheaded fireworm Sparganothis fruitworm Armyworm (e.g. false) Blackvine weevil**	Four (1/3 lb. WSP)	Use sufficient water to give thorough coverage with ground or sprinkler equipment. For Black Vine Weevils, make application night after peak adult emergence and prior to egg laying. If applied through an irrigation system, refer to "Sprinkler Application To Cranberries" statement on product label for specific application instructions and requirements. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or broadleaf weeds. Do not apply this product or allow it to drift to blooming small fruits or broadleaf weeds: apply as a pre-bloom or post-bloom spray only	75

*Pheromone traps are available for some insects which are very helpful in predicting egg hatch and early larval stages. Frequent scouting is also advised. Consult local Extension Service personnel and/or Consultants for information on application timings based on trap catches and other Integrated Pest Management (IPM) procedures.

**Suppression only.

RESTRICTIONS / PRECAUTIONS

- Do not apply less than 75 days before harvest.
- Do not apply more than 1.0 lb. of active ingredient or more than one application per acre per crop cycle.
- Do not apply from start of bloom until all berries set.
- Do not apply using aerial equipment.
- Tank mix post-bloom application of Orthene with another organophosphate insecticide product, such as Diazinon, is at the risk of the user, applicator and/or applicator advisor.

WSDA Container Disposal Guidance: Completely empty the contents of the pesticide container into application equipment and apply to a labeled crop or site. The empty container may be disposed of in a sanitary landfill. Burning is not a legal method of container disposal in Washington.

This label for *Orthene* 75 S Soluble Powder (Water Soluble Packets) expires and must not be distributed or used in accordance with this SLN registration after December 31, 2012.

FIFRA § 24(c) Registrant: Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596-8025

Orthene is a registered trademark of OMS Investments, Inc.