



Bayer CropScience LP  
P.O. Box 12014  
2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)  
<http://www.bayercropscience.com>

ROVRAL® brand 4 Flowable Fungicide  
For Use on Crucifer Crops Grown for Seed  
EPA Reg. No. 264-482  
EPA SLN. No. WA-960027

## 24(c) Supplemental Label

### FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

This label for ROVRAL® brand 4 Flowable Fungicide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2010.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label. This product may be applied through irrigation systems.

CROP	INSECT	RATE	REMARKS
Crucifer crops	Alternaria leaf and pod blight, Sclerotinia white rot, Black leg (Phoma lingam)	Use 2 to 4 pints of product per acre per application. The higher rate should be used when severe disease conditions exist.	Crucifer crops that are grown for seed and listed in Washington State rules WAC 16-228-1270 only (including, but not limited to, broccoli, Brussels sprouts, cabbage, cauliflower, kohlrabi, kale, radish, rape, rutabaga and turnip).

#### Dilution Rate and Method of Application:

Apply in 20 to 100 gallons of water per acre by ground. Apply in a minimum of 10 gallons of water per acre by air. Addition of a suitable resin-based surfactant is recommended according to dilution rates on the surfactant label. Thorough coverage is necessary for effective control.

Frequency and Timing of Application: Apply Rovral 4F when Brassica seedlings are transplanted in late summer or early fall, to reduce the risk of black mold during cool and moist conditions. Applications should also be made at full bloom, at pod set and just prior to harvest if favorable conditions exist.

#### RESTRICTIONS:

- All Crucifer crop seed screenings shall be disposed of in such a way that they cannot be distributed or used for food or feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the WSDA forthwith upon request. Disposal records shall consist of documentation from a controlled dumpsite, incinerator, or other equivalent disposal site and shall show the lot numbers, amount of material disposed of, its grower(s), and the date of disposal.
- No portion of the Crucifer seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
- Crucifer seed crop shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.
- Crucifer seed crop may not be distributed for human consumption or animal feed.
- If applying this product adjacent to a water body such as a lake, reservoir, river, permanent stream, marsh or natural pond, estuary, or commercial fishpond, there must be at least a 25-foot buffer between the water body and the point of application.

This pesticide is toxic to invertebrates. ROVRAL® brand 4 Flowable Fungicide should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program website at <http://agr.wa.gov/PestFert/EnvResources/EndangSpecies.htm>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the

pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the internet from WSU at <http://pep.wsu.edu/waste/wd.html> or from WSDA at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

#### FUNGICIDE RESISTANCE STATEMENT

ROVRAL® brand 4 Flowable Fungicide is a dicarboximide fungicide. Repeated use of dicarboximide fungicides may result in resistance to ROVRAL®.

NOTE: Consult the product label for all handling directions.

ROVRAL is a registered trademark of Bayer.