

REGISTRATION FOR SPECIAL LOCAL NEED
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

GOWAN CRYOLITE BAIT

EPA Reg. No. 10163-225
EPA SLN NO.: WA-980001

KEEP OUT OF REACH OF CHILDREN CAUTION

This label for Gowan Cryolite Bait expires and must not be distributed or used in accordance with this SLN registration after December 31, 2012

DIRECTIONS FOR USE

- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of pesticide application.
- All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

GENERAL USE DIRECTIONS AND PRECAUTIONS

GOWAN CRYOLITE BAIT is recommended as part of an Integrated Pest Management program for control of insects listed. Apply by ground equipment only. Product should be applied uniformly around the base of plants.

CROP RECOMMENDATIONS

CROP	PEST	PRODUCT RATE (LBS) / ACRE	APPLICATION DIRECTIONS
BLUEBERRIES, RASPBERRIES, STRAWBERRIES	Black Vine Weevil, Rough Strawberry Root Weevil, Strawberry Root Weevil	20-40	Based on pest pressure, a second application may be made 10-14 days later.

RESTRICTIONS / PRECAUTIONS: This product is formulated for application by ground equipment only. Do not use for control of obscure root weevil. Do not apply within 3 days of harvest. Do not use GOWAN CRYOLITE BAIT in combination with lime or compounds containing free lime. Do not apply under conditions involving possible drift to food, forage or other planting that might be damaged, or the crops thereof rendered unfit for sale, use or consumption. Cryolite is a stomach poison. Sufficient amounts of bait must be ingested by insects in order to achieve control. To minimize plant damage, make application at first indication of insect pressure. Application equipment should be properly calibrated to ensure a uniform application rate and complete coverage. Significant rain and/or irrigation immediately following application may reduce bait effectiveness. Do not apply under very dry soil conditions.

WSDA Container Disposal Guidance: Completely empty the contents of the pesticide container into application equipment, and apply to a labeled crop or site. The empty container may be disposed of in a sanitary landfill. Burning is not a legal method of container disposal in Washington.

24(c) Registrant:
Gowan Company
P.O. box 5569
Yuma, AZ 85366



Rev. 7/15/08