

# DUPONT CROP PROTECTION

## SPECIAL LOCAL NEED

### 24(C)LABELING

#### DUPONT™ CURZATE® 60 DF FUNGICIDE

FOR THE CONTROL OF DOWNY MILDEW IN  
SPINACH, CABBAGE, SWISS CHARD, AND TABLE  
BEET GROWN FOR SEED IN THE STATE OF  
WASHINGTON

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

## DUPONT™ CURZATE® 60 DF FUNGICIDE

EPA REG. NO. 352-592  
EPA SLN NO. WA-990021

### FOR CONTROL OF DOWNY MILDEW IN SPINACH, CABBAGE, SWISS CHARD AND TABLE BEET GROWN FOR SEED

#### DIRECTIONS FOR USE

It is a violation of federal law to use this producer in a manner inconsistent with its labeling.

DuPont CURZATE® 60 DF fungicide may be applied to spinach, cabbage, Swiss chard and table beet grown for seed production only for the control of downy mildew. Consult the CURZATE® 60 DF EPA-registered label for additional requirements, restrictions and precautions.

#### IMPORTANT

Before using CURZATE® 60 DF fungicide, follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA-registered label.

#### HOW TO USE

##### Use Rate, tank mixes and timing

- Apply CURZATE® 60 DF fungicide at 3.2 ounces per acre.
- Use CURZATE® 60 DF only in combination with another registered protectant fungicide, such as Manzate® 75DF, or chlorothalonil.
- Applications can be made prior to or after transplanting.
- Initial applications should start when local conditions indicate that downy mildew infection is imminent.
- Make applications on a 5-7 day interval as long as disease conditions exist.
- Do not exceed 5 applications of CURZATE® 60 DF per 12 month period.

#### Application information

- Do not apply by airplane.
- Do not apply this product through any type of irrigation system.
- Apply CURZATE® 60 DF in sufficient water to obtain thorough spray coverage of the crop.
- For conventional ground application equipment, including tractor-mounted sprayers, boom or backpack sprayers, use a minimum of 20 gals/acre.
- For hard-to-wet foliage, a spray adjuvant may be used to help increase spray coverage and retention.

#### USE PRECAUTIONS AND RESTRICTIONS

- For use only on crops grown for seed.
- All cabbage, spinach, Swiss chard and table beet seed screenings shall be disposed of in such a way that they cannot be distributed or used for food or feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the WSDA forthwith upon request. Disposal records shall consist of documentation from a controlled dumpsite, incinerator, or other equivalent disposal site and shall show the lot numbers, amount of material disposed of, its grower(s), and the date of disposal.
- No portion of the cabbage, spinach, Swiss chard and table beet seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.

For product information call 1-888-6-DUPONT

© 2005 E. I. du Pont de Nemours and Company, Agricultural Products, Wilmington, Delaware 19898

- Cabbage, spinach, Swiss chard and table beet seed shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.
- Cabbage, spinach, Swiss chard and table beet seed may not be distributed for human consumption or animal feed.
- Do not replant treated fields to any crop within 30 days after last CURZATE® 60 DF treatment.

#### **Environmental Hazards**

This product is toxic to fish and aquatic invertebrates. DuPont™ Curzate® 60DF fungicide should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program website at <http://agr.wa.gov/PestFert/EnvResources/EndangSpecies.htm>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed threatened or endangered species. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

#### **IMPORTANT**

**BEFORE USING CURZATE® 60 DF FUNGICIDE, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.**

#### **WSDA Container Disposal Guidance:**

Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the internet from WSU at <http://pep.wsu.edu/waste/wd.html> or from WSDA at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

Read the Limitation of Warranty and Liability on the section 3 Federal product label before buying or using this product. If terms are not acceptable, return the unopened package at once to seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

This labeling must be in the possession of the user at the time of pesticide application.

**This label for DuPont CURZATE® 60 DF fungicide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2009.**

Manzate® is a registered trademark of DuPont.

R-379 010605

